# **Plants of South Eastern New South Wales**




Flower. Photographer Don Loarie, California, USA



Flower. Photographer Don Loarie, California, USA



Leaves and seeds. Photographer Ori Fragman Sapir, Israel



Line drawings. d. petal; petal base and nectary; leaf; seed. BMMayfield, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

### Common name

Sharp Buttercup, Burr Buttercup, Prickle-fruit Buttercup

### **Family**

Ranunculaceae

#### Where found

Lawns, pastures, gardens, roadsides, disturbed sites, stream banks, moist gullies, heavy waterlogged soils, and bogs. Widespread. No records from Kosciuszko National Park.

## **Notes**

Introduced annual herb, to 0.5 m high, stoloniferous, hairless, sparsely hairy or slightly rough. Leaves basal and alternating up the stems, 1–5 cm long, 10-50 mm wide, entire or 3-5 lobed, margins scalloped or toothed. Upper stem leaves more or less wedge shaped, with 3 narrow, entire or toothed, lobes. Flowers yellow, glossy, with 5 petals each 6–10 mm long. Sepals strongly turned down. Flowers in loose 1-10-flowered clusters. Flowers Aug.—Nov. Seeds 8–25, brown, broadly oval, flattened, 5–8 mm long, margin strongly keeled and grooved, faces with stout wart-based tapering spines or rarely almost smooth, beak stout, slightly curved down to almost straight, 1.5–4 mm long.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Ranunculus~muricatus (accessed 4 February, 2021)

Author: Betty Wood.

 $This identification \ key \ and \ fact \ sheets \ are \ available \ as \ a \ free \ mobile \ application:$ 





