
Plants of South Eastern New South Wales



Flowers and leaves. Photographer Don Wood, SE Forests National Park



Rants. Australian Rant Image Index, photographer Murray Fagg, Namadgi National Park, ACT



Line drawings. b. plant; leaf with detail of leaf stalk; petal base and nectary; seed. BM/Mayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Bog buttercup

Family

Ranunculaceae

Where found

Forest, woodland, alpine **herbfields**, wet **grassland**, and swamps and alpine bogs, predominantly at 800-2000 m altitude. Kosciuszko National Park, the mountains to the north, ACT, tablelands, and ranges.

Notes

Perennial herb to 0.25 m high, hairy, rarely almost hairless. Leaves basal, 0.5–5 cm long, 5–25 mm wide, hairy with fine spreading hairs, compound, with 3–7 leaflets which are further lobed, or or with conspicuous rounded teeth. Flowers yellow, 5–18 mm in diameter, with 5-8 petals. Sepals spreading, slightly rough. Flowering stems unbranched, or branched with up to 5 flowers. Flowers spring-summer. Seeds usually 10–40, 1.5–3 mm long, approximately elliptical, flattish, the sides mildly convex, smooth, tapering gradually into a stout beak 0.5–1 mm long, usually with a hooked or curved down tip.

Hybridises with Ranunculus dissectifolius and Ranunculus graniticola.

Some small specimens from Kosciuszko National Park, and the Brindabella Range, ACT, appear to be intermediate between Ranunculus pimpinellifolius and Ranunculuus productus.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Ranunculus~pimpinellifolius (accessed 4 February, 2021) Author: Betty Wood.

6 ۲ This identification key and fact sheets are available as a free mobile application:



Android edition



Creative Commons Attribution 3.0 Australia (CC BY)