Plants of South Eastern New South Wales

Common name

A Buttercup

Family

Ranunculaceae

Where found

Wet montane and subalpine grassland at about 1500-1800 m altitude. Kosciuszko National Park.

Notes

Perennial herb. Flowering stems mostly 0.01–0.04 m long at flowering, elongating to 0.45 m in seed. Stems hairy with short appressed hairs. Leaves mostly basal, 0.7–3.5 cm long, 7–20 mm wide, hairy with short appressed hairs, compound, with 3–7 oval leaflets which are 3-lobed and further lobed or toothed. Flowers pale yellow, with 5 petals, each 1.5-2.5 mm long. Flowering stems 1-flowered. Sepals spreading, rarely slightly turned down. Flowering: Dec.-Mar. Seeds 25–45, lens-shaped, elliptical to almost round, 0.3–2 mm long, not strongly compressed, side faces smooth; beak 0.3-0.8 mm long, slender, strongly curved down.

Some small specimens from Kosciuszko National Park, and the Brindabella Range, ACT, appear to be **intermediate** between **Ranunculus pimpinellifolius** and **Ranunculuus productus**.

PlantNET description with line drawings and photo: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Ranunculus~productus (accessed 4 February, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:







Creative Commons Attribution 3.0 Australia (CC BY)