

Plants of South Eastern New South Wales




Flowering stem. Photographer Don Wood, west of Nowra



Fruit and leaves. Photographer Daniel Ohlsen, © 2021 Daniel Ohlsen



Flowers and leaves. Photographer Neville Walsh, © 2021 Royal Botanic Gardens Board, Melbourne, Vic



Line drawing. b. fruiting stem. EMMayfield, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Rose-leaf Bramble, Native Raspberry

Family

Rosaceae

Where found

Forest. Coast, ranges, and Southern Highlands.

Notes

Shrub to 2 m high, trailing or **scrambling**. Prickles 2-15 mm long, scattered on the stems, leaf stalks. **Rachis** and lower midveins of the leaves, and flower stalks. Fruit fleshy. Stems sticky hairy, sometimes almost hairless. Leaves **alternating** up the stems, 5-17 cm long, **compound**, usually with 3-7 leaflets each 3-8 cm long, 10-35 mm wide, margins irregularly and often deeply toothed, both surfaces with **glandular** hairs or bearing small resinous globules, **main vein** usually more densely hairy. Leaves **simple** near the flower clusters. Flowers white or pale pink, with 5 **petals** each 5-13 mm long, broadly elliptic to almost round. Flowers single or in short branched clusters. Fruit red, oval to oval, 15-20 mm long.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Rubus-rosifolius> (accessed 5 February, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)