
# **Plants of South Eastern New South Wales**



Flowering stem. Photographer Don Wood, Coolungubra State Forest near Bombala



Flowering plant. Australian Flant Image Index, photographer Murray Fagg, near Willis, NSW



Line drawings. c. flowering branch; d. flower. BM Mayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

## Common name

Soapwort, Bouncing Bet, Common Soapwort

#### **Family**

Caryophyllaceae

### Where found

Weed of roadsides, waste ground and along streams. Widespread, but rarely coastal except in the Sydney area.

#### **Notes**

Introduced perennial herb to 1 m high, occasionally prostrate, with stout branching rhizomes and long stolons. Stems hairless or with sparse hairs. Leaves opposite each other, 2.5–10 cm long, 6–30 mm wide, conspicuously 3-veined, lower surface covered with minute blisters, tips pointed. Flowers white or pink to pinkish mauve, tubular, with 5 lobes, each about 10 mm long. Flowers in tight clusters of 6–25 flowers. Flowering: winter to summer.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Saponaria~officinalis (accessed 6 February, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





