# **Plants of South Eastern New South Wales**




Flower head. Photographer Sue Guymer, east of Melbourne



Flow ering plant. Photographer Sue Guymer, east of Melbourne



Seeding plant. Photographer Russell Best, north of Sunbury, Vic



Line drawings. a. plant; leaf margin detail; seed. E Mayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

### Common name

A Solengyne

## **Family**

Asteraceae

### Where found

Forest, woodland, and grassland, often persistent in lawns. Widespread. Rare on the coast and Western Slopes.

#### **Notes**

Perennial herb to 0.11 m high. Flower stalks robust, about 0.5-1 mm in diameter. Stems hairy to rough. Leaves basal, 1–11 cm long, 5–22 mm wide, surfaces sparsely to densely hairy, margins with 1–15 teeth, often confined to the upper half of the leaf, occasionally scalloped or wavy, tips blunt. Flower heads 3–7 mm in diameter, hemispherical. 'Petals' pale to magenta or purplish, minute. Flower heads single. Flowering: spring—autumn. Bracts appressed to the flower heads erect at the seeding stage.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Solenogyne~gunnii (accessed 7 February, 2021)

Author: Betty Wood.

 $This identification \ key \ and \ fact \ sheets \ are \ available \ as \ a \ free \ mobile \ application:$ 





