
# **Plants of South Eastern New South Wales**



Hower cluster. Photographer Jackie Mles



Flow ering stem. Photographer Don Wood, Booderee National Park, Jervis Bay Territory



Line drawings. e. flowering branch; f. flower. BM Mayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

### Common name

Pink swamp heath

## **Family**

Ericaceae

### Where found

Swampy shrubland and wet heath and sedgeland. Coast and ranges.

#### **Notes**

Shrub to 2 m high, hairless. Stems smooth. Leaves alternating up the stems, overlapping each other, 0.3–2 cm long, 2–10 mm wide, concave; tips gradually tapering to a point, and with a mucro to 0.7 mm long, margins entire, or fringed on the stem-clasping part of the leaf, bases completely sheathing the stems. Flowers pink, purplish, or white, tubular, the tube 0.5–1.3 mm long, with 5 spreading lobes each 3–5 mm long. Flowers in clusters of 3–20, often appearing to form many-flowered clusters. Flowers most of the year.

Family was Epacridaceae.

Protected NSW.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Sprengelia~incarnata (accessed 7 February, 2021)

Author: Betty Wood.

 $This \, identification \, key \, and \, fact \, sheets \, are \, available \, as \, a \, \, free \, mobile \, application: \, \\$ 





