# **Plants of South Eastern New South Wales**



Male flowers and leaves. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Fermale flowers and leaves. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT



Fruit and leaves. Australian Flant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra, ACT

## Common name

Brush pepperbush, Dorrigo pepper, Scrub pepperbush

## **Family**

Winteraceae

### Where found

In or near rainforest, and wet gullies. Coast, ranges, and the mountainous parts of the tablelands, north of Narooma.

#### **Notes**

Shrub or tree to 5 m high. Fruit fleshy. Stems hairless. Leaves without a distinctive scent when rubbed, but with a peppery taste. Leaves alternating up the stems, whorled, or clustered, 4–20 cm long, 13–55 mm wide, bases usually auriculate, surfaces hairless, green and glossy, tips gradually tapering to a point. Male and female flowers on different plants. Flowers white to cream or yellowish, with 2-4 petals, each 5–13 mm long. Flowers in clusters of 1-16 flowers. Flowering: August–November. Fruit usually purplish, sometimes whitish mottled with red, more or less oval, about 12–20 mm long.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Tasmannia~insipida (accessed 9 February, 2021)

Author: Betty Wood.

 $This \, identification \, key \, and \, fact \, sheets \, are \, available \, as \, a \, \, free \, mobile \, application: \, \\$ 







Creative Commons Attribution 3.0 Australia (CC BY)