Plants of South Eastern New South Wales



Flowers and leaves. Australian Flant Image Index. photographer Murray Fagg, Walcha



Seed case. Photographer Robert Hille, unknown place



Line drawings. a. leaf; flower head; seeding calyx. M Mbir, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Red clover

Family

Fabaceae

Where found

Grassy areas and disturbed sites. Widespread.

Notes

Introduced perennial herb to 1 m tall or sprawling. Calyx teeth blunt. Stems moderately hairy to hairless. Leaves alternating up the stems, compound, with 3 leaflets, each 1.5-6 cm long, 7-35 mm wide, appressed-hairy on both surfaces or on just the lower surface, surfaces sometimes spotted, tips pointed, blunt, or rounded with a small notch, margins virtually entire, stipules fused to the stalks, membranous, veined, and hairy or becoming hairless. Flower heads globular or oval, 14-30 mm long, with more than 100 flowers. Flower heads single or paired. Individual flowers pink or pinkish red to purple, 14-18 mm long, pea shaped, with 5 petals, 2 joined together to form the keel. Flowers mainly Oct.—May.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Trifolium~pratense (accessed 3 May 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





