
Plants of South Eastern New South Wales



Flowering stems. Australian Flant Image Index, photographer Murray Fagg, Mulligans Flat Nature Reserve, ACT



Seed case and leaves. Photographer Zoya Akulova, California, USA



Line drawings. f. leaf; flower head; seeding calyx. M Moir, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Knotted clover

Family

Fabaceae

Where found

Grassy areas and disturbed sites. Widespread. Rare on the coast and ranges.

Notes

Introduced annual herb to 0.6 m tall or spreading. Calyx teeth spiny, erect or slightly spreading at the seeding stage. Stems sparsely to densely hairy. Leaves alternating up the stems, compound, with 3 leaflets, each 0.3-2.5 cm long, 1.5-13 mm wide, appressed-hairy on both surfaces, lateral veins mostly straight near the margins, upper margin with minute teeth, tips with a minute mucro, and blunt to rounded with a small notch; stipules submembranous, appressed-hairy, green-veined, fused to the stalks for up to half their length. Flower heads oval to cylindrical, 6-20 mm long, 8-15 mm wide, 20-40-flowered. Individual flowers pink, 5-7 mm long, pea shaped, with 5 petals, 2 joined together to form the keel, persistent at the seeding stage. Flowers mainly Oct.—Jan. Calyces surrounding the individual pods at the seeding stage.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Trifolium~striatum (accessed 3 May 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





