Plants of South Eastern New South Wales




Flower head and leaves. Australian Flant Image Index, photographer Kevin Thiele, WA



Seeding stems. Photographer A.J. Brown, © 2021 Royal Botanic Gardens Board, Melbourne, Vic



Seeding plant. Photographer Nimal G. Karunajeewa, © 2021 Royal Botanic Gardens Board, Melbourne, Vic



Line drawings. h. leaf; flower head; seeding calyx. M Moir, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Woolly Clover

Family

Fabaceae

Where found

Naturalised. Sydney area, Canberra, and Western Slopes. Occasionally elsewhere.

Notes

Introduced annual herb to 0.6 m high or sprawling to prostrate. Calyx teeth blunt. Stems cylindrical or furrowed, and hairless to sparsely hairy. Leaves alternating up the stems, compound, with 3 leaflets, each 0.4–1.5 cm long, 2–10 mm wide, wholly hairless or sparsely hairy on the lower surface, margins toothed, tips rounded, squared off, or rounded with a small notch; stipules membranous, joined together in the lower half. Flower heads many-flowered, 6–12 mm long, 5–8 mm wide. Individual flowers pink, 3–6 mm long, pea shaped, with 5 petals, 2 joined together to form the keel, falling early. Calyces of individual flowers at first about 3.5 mm long, later much inflated at the pod stage, and woolly-hairy to hairless. Flowers Sep.–Nov.

Varieties not recognised in NSW.

PlantNET description (as *Trifolium tomentosum*): http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Trifolium~tomentosum (accessed 3 May 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:





