


Plants of South Eastern New South Wales



Flowers. Australian Plant Image Index, photographer Murray Fagg, Australian National Botanic Gardens, Canberra



Flowering stem Photographer Don Wood, Boyne State Forest, north east of Nelligen



Line drawings. h. flowering branch; seed case; seed. E Mayfield, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Bearded tylophora

Family

Apocynaceae

Where found

Forest and along streams. Coast, ranges, and occasionally on the tablelands.

Notes

Slender twiner, trailing plant, or vine, to 3 m tall. Stems more or less hairless. Leaves opposite each other, 2.5–7.5 cm long, 10–40 mm wide, thin, both surfaces more or less hairless, tips pointed. Flowers 5–9 mm in diameter, dark red or dark brown to reddish purple, tubular, with 5 **spreading** lobes, in 3–5-flowered short clusters in groups of 1–3. Flowers spring–summer.

Family was Asclepiadaceae.

PlantNET description: <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Tylophora~barbata> (accessed 26 March, 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)