


Plants of South Eastern New South Wales



Flowering plant. Photographer Jackie Miles, Kosciuszko National Park



Flowering stems. Photographer Don Wood, South East Forests National Park



Line drawings. plant, flower. S Clark, National Herbarium of Victoria, ©2021 Royal Botanic Gardens Board

Common name

Tasmanian bladderwort

Family

Lentibulariaceae

Where found

Shallow water and wet soil near streams and ponds. Kosciuszko National Park. Ranges and eastern edge of the tablelands south of Bombala

Notes

Perennial herb, **stoloniferous**. Flower stalks to 0.15 m high. Insect traps moderately numerous on the 'roots'. Leaves numerous, in **rosettes** and on the stems, to about 3 cm long, about 1 mm wide. Flowers mostly dark violet, 6–10 mm long, upper lip round to oval, tip notched, the lower lip much larger, transversely elliptic, with 3 conspicuous yellow central ridges flanked by several less conspicuous violet ridges. Flowers single, occasionally paired. Flowers December–March.

Vulnerable Vic. Listed in the Flora and Fauna Guarantee Act, Vic.

VICFLORA description: <https://vicflora.rbg.vic.gov.au/flora/taxon/30c4d90b-0685-4dd3-b6e4-0694b3713063> (accessed 3 May 2021)

Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)