Plants of South Eastern New South Wales



Flowering stems. Photographer Don Wood, Bungonia State Conservation Reserve



Leafy stemwith flower. Photographer Jackie Mles



Line drawings. g. flow ering and seeding stem; seed case; stem detail. S Oark, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Trailing Speedwell, Creeping Speedwell

Family

Plantaginaceae

Where found

Forest, grassland, and a weed in lawns and gardens. Widespread.

Notes

Creeping perennial herb to 0.1 m high, with stolons to about 1 m long. Stems hairy with very short hairs, denser in the longitudinal bands, and longer and denser at the nodes. Leaves opposite each other, mostly 0.8-2 cm long, 5-16 mm wide, margins with 3-8 pairs of irregular teeth, bases squared off to slightly cordate, tips more or less pointed. Flowers mauve, 2-3.5 mm long, with a short tube and 4 spreading lobes. Stamens 2. Flowers in clusters of 3–10 flowers. Flowering: spring-summer.

Family was Scrophulariaceae.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfI.pl? page=nswfl&lvI=sp&name=Veronica~plebeia (accessed 11 February, 2021)

Author: Betty Wood.

Θ

This identification key and fact sheets are available as a free mobile application:



Android edition iOS edition eń.

Creative Commons Attribution 3.0 Australia (CC BY)