Plants of South Eastern New South Wales



Flowers. Photographer Richard Hartland, Bogong High Plains, Vic



Flowering stem Australian Plant Image Index, photograher AN Schmidt-Lebuhn, Tas



Flow ering plants. Australian Plant Image Index, photograher Colin Totterdell, Kosciuszko National Park



Line drawings. a. flowering plant; flower; seed case. E Mayfield, National Herbarium of Victoria, © 2021 Royal Botanic Gardens Board

Common name

Waxy bluebell

Family

Campanulaceae

Where found

Moist sites in heath, grassland, along streams, and around and in swamps at high montane to alpine altitudes. Ranges, tablelands, ACT, the mountains to the west, and Kosciuszko National Park.

Notes

Perennial herb to 1.15 m high, with spreading rhizomes. Stems at first sprawling, finally erect, mostly unbranched, occasionally 1or 2-branched, hairless or sometimes sparsely rough-hairy towards the base. Leaves alternating up the stems, 0.5–5 cm long, 1–10 mm wide, usually hairless, margins usually flat, sometimes more or less wavy but less so than Wahlenbergia gloriosa, often toothed, tips more or less blunt to pointed. Flowers sky blue to pale violet, occasionally white, often whitish outside, with a bell-shaped tube 3-9 mm long, and 5 lobes 10-23 mm long. Corolla lobes with long drawn out points. Stigma usually with 3 lobes, occasionally 2. Flowers single or 2-3 per stem, nodding in bud. Flowering: summer-autumn.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Wahlenbergia~ceracea (accessed 8 February, 2021)

Author: Betty Wood

This identification key and fact sheets are available as a free mobile application:





