Plants of South Eastern New South Wales



Flowers. Photographer Richard Hartland, Mt Stirling, Vic



Flowering plant. Australian Plant Image Index, photographer Murray Fagg, Namadgi National Park, ACT



Photographer Don Wood, Namadgi National Park, ACT



Line drawings. a. flowering branch; seed case. E Mayfield, National Herbariumof Victoria, © 2021 Royal Botanic Gardens Board

Common name

Royal bluebell

Family

Campanulaceae

Where found

Forest, woodland, alpine herbfields and grassland, roadsides, rocky sites, and moist sites, at high altitudes. South and east of the Hume Highway. Kosciuszko National Park, the mountains to the north, ACT, tablelands and ranges.

Notes

Perennial herb to 0.4 m high, with spreading rhizomes. Stems sprawling to erect, mostly unbranched, usually hairy to densely hairy towards the base, sometimes hairless. Leaves often mostly at the base of the plant, stem leaves mostly opposite each other, sometimes the upper leaves, rarely all leaves, alternating up the stems, 0.4–3.5 cm long, 1–15 mm wide, margins usually wavy, entire, or with small hardened teeth, surfaces hairless or the lower leaves hairy, tips blunt to pointed. Flowers purple to purplish violet, or deep blue, with a bell-shaped tube 2.5–8.5 mm long, and five lobes 11–25 mm long. Stigma usually with 2 lobes, sometimes 3. Flowers single. Flowering: summer-autumn.

Hybrids between Wahlenbergia gloriosa and Wahlenbergia stricta subsp. stricta have been recorded.

All native plants on unleased land in the ACT are protected.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl? page=nswfl&lvl=sp&name=Wahlenbergia~gloriosa (accessed 8 February, 2021) Author: Betty Wood.

This identification key and fact sheets are available as a free mobile application:



Android edition



iOS edition

Creative Commons Attribution 3.0 Australia (CC BY)