Plants of South Eastern New South Wales



Flowers. Australian Plant Image Index, photographer Murray Fagg, Nadgigomar north of Braidwood.

Common name

A Grass-tree

Family

Xanthorrhoeaceae

Where found

Dry forest and heath, often in seasonally waterlogged sites. Mainly coastal.

Notes

Shrub-like perennial herb. Trunk absent. Scape below the spike 0.5–2.5 m long, about 10 mm in diameter; flower spike 0.5–0.9 m long, 15-30 mm in diameter. Leaves basal, forming a spreading to almost erect tuft. Leaves 3-6 mm wide, 1.5-2 mm thick, bluegreen, glaucous, almost a wide triangle in cross section, the long side almost flat or concave. Individual flowers with 6 'petals' in two rows, outer 'petals' papery or more or less membranous, inner 'petals' membranous. The whole spike light brown when flowering. Bracts surrounding the flowers obscure. Bracts between the flowers densely hairy with pale hairs, appearing velvety. Flowering: October-December

Protected NSW.

PlantNET description: http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfI.pl? page=nswfl&lvl=sp&name=Xanthorrhoea~concava (accessed 12 February, 2021)

Author: Betty Wood.



This identification key and fact sheets are available as a free mobile application:





Creative Commons Attribution 3.0 Australia (CC BY)



Flow ering plant. Photographer Don Wood, Boyne State Forest north of Batemans Bay