



Pacific Pests and Pathogens - Fact Sheets

<https://apps.lucidcentral.org/ppp/>

Banana bunchy top (121)



Photo 1. Banana plants of different ages showing symptoms of *Banana bunchy top virus* disease. Notice the stunted plants and the colour of the leaves, which are pale yellow at the margins.



Photo 2. Symptoms of *Banana bunchy top virus*. Notice the leaves are upright, stunted and tend to cluster in the throat of the plant; leaves like this are said to be "choked", and give the bunchy top symptom.



Photo 3. Underside of a banana leaf showing veins that have dark and light areas (dots and dashes) along them. The veins near the midrib bend down, "hooking" into it.



Photo 4. Colony of the banana aphid, *Pentalonia nigronervosa*. Note the dark colour of the adults and the nymphs. The dark veins of the winged adults can just be seen on the insect at the top left corner.

Common Name

Banana bunchy top

Scientific Name

Banana bunchy top nanavirus. The abbreviation is BBTV.

Distribution

Widespread. Asia, Africa, North America (Hawaii), Oceania. It is recorded from American Samoa, Australia (restricted), Fiji, Guam, New Caledonia, Samoa, Tonga, Tuvalu, Wallis and Futuna.

Hosts

Banana and others in the genus *Musa*.

Symptoms & Life Cycle

Plants with banana bunchy top (BBT) have narrow, progressively shorter leaves with little distance between them, hence the name

'bunchy top' (Photos 1&2). Leaf edges often roll upwards with yellow margins. The most characteristic symptoms are short, dark-green dots and dashes along the minor leaf veins, so-called 'Morse-code' patterns, which form hooks curving downwards near the edge of the midrib (Photo 3). Hold the leaf up to the light and view from the underside.

The virus is spread by an aphid, *Pentalonia nigronervosa* (Photo 4). It takes more than 4 hours of feeding on a diseased plant before aphids become infected with the virus. But once infected, the aphids continue spreading the virus until they die. Aphids live for 15-20 days. The virus is not transmitted on tools or through the soil (**see also Fact Sheet no. 103**).

Impact

BBT is caused by a virus. In the past, the disease has devastated banana industries in Australia, Fiji, Samoa and Tonga. Today, it is causing much damage in Pakistan and is spreading in the DR Congo. Plants that are infected when young rarely produce a fruit bunch. When diseased suckers are planted they become severely stunted and they do not produce fruit. Plants infected at a later growth stage may produce a distorted bunch. In Pacific island countries, the failure to select healthy planting material has resulted in plantations showing 20% of plants with BBT in the first year, and major losses within 2-3 years.

Detection & inspection

Look for plants with stunted, short, bunched leaves, with the "Morse-code" patterns (dot-dash) symptom. Yellowing of the leaf margins also suggests BBT, though this can be caused by other problems. There are commercial ELISA kits to detect the disease. Molecular techniques using PCR (polymerase chain reaction) are more sensitive.

Management

QUARANTINE

Banana plants should only be transferred between countries as tissue cultures tested to ensure that they are free of all known banana diseases. Plants of many varieties of bananas, tested to internationally agreed standards, can be obtained from the INIBAP transit centre in Belgium, and from IITA, Nigeria. The FAO/IBPGR *Technical Guidelines for the Safe Movement of Germplasm*. No. 15. Musa. 2nd Edition, gives details of the methods to use (http://www.bioversityinternational.org/uploads/tx_news/Musa_spp._2nd_edition__502.pdf).

NATURAL ENEMIES

In Tonga, *Aphidius colemani*, a parasitic wasp, was released for biological control of the aphid, but results were disappointing.

CULTURAL CONTROL

Key actions in managing BBT are: (i) the monitoring of banana plantations for BBT symptoms; and (ii) rapid removal of diseased plants. If the disease is found in the field, do the following:

- Before destruction, spray diseased plants with soap, oils, kerosene or insecticide to kill all aphids (see below).
- After killing the aphids, you have three choices:
 - Method 1: Within 48 hours, dig out the stool, including the main plants and all suckers. Chop the plants, including the corms, into small pieces. Burn or bury.
 - Method 2: Inject the diseased plant with herbicide (e.g., glyphosate, 1 part to 5 parts water). Inspect the plants to make sure they are dead. If not, retreat.
 - Method 3: Scoop out the middle of the stem and pour kerosene into the well. Inspect the plants to make sure they are dead. If not, retreat.

RESISTANT VARIETIES

All varieties of banana are susceptible, although some may take longer to develop symptoms. Some varieties may escape infection because they are not as attractive to aphids as other varieties.

CHEMICAL CONTROL

The choice of chemicals for killing aphids on diseased plants, before removal from the soil, depends on whether they are grown for household use or for sale.

- *Banana for home use*: Use soap solution, white oil (vegetable oil) or horticultural oil (petroleum oil) (**see Fact Sheet no. 56**). Kerosene can also be used as mentioned above. Aim to spray the "throat", the V-shaped area where the leaves meet and where

aphids hide.

- *Commercial plantations*: The following have been recommended for Pacific island countries: dimethoate (400 g/L), used at 75 ml/100L; diazinon (200 g/L), used at 1.5 ml/L; acephate (75% WP), used at 1.3 g/L. Note, in Australia, the use of dimethoate is restricted, and the use of the chemical remains under review. Diazinon use is also under review.
- Synthetic pyrethroids are likely to be effective.

AUTHOR Grahame Jackson

Information from CABI (2015) *Banana bunchy top virus* (bunchy top of banana) Crop Protection Compendium. (www.cabi.org/cpc). Photo 1 Richard Markham. ACIAR, Canberra.

Produced with support from the Australian Centre for International Agricultural Research under project PC/2010/090: *Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This fact sheet is a part of the app *Pacific Pests and Pathogens*

The mobile application is available from the Google Play Store and Apple iTunes.



Copyright © 2019. All rights reserved.