



Pacific Pests and Pathogens - Fact Sheets

<https://apps.lucidcentral.org/ppp/>

Tobacco flea beetle (355)



Photo 1. Adult tobacco flea beetle, *Epitrix hirtipennis*.
Note, the long antennae.



Photo 2. Adult tobacco flea beetle, *Epitrix hirtipennis*.
Side view.



Photo 3. Tobacco flea beetle, *Epitrix hirtipennis*,
feeding on tomato



Photo 4. Adult tobacco flea beetle, *Epitrix hirtipennis*,
and holes eaten in a tomato leaf.



Photo 5. Severe damage on eggplant by the tobacco flea
beetle, *Epitrix hirtipennis*.



Photo 6. Severe damage on eggplant by the tobacco
flea beetle, *Epitrix hirtipennis*.



Photo 7. Large swarm of tobacco flea beetle, *Epitrix hirtipennis*, on eggplant.



Photo 8. Irregular patches on leaf of tobacco where surface layers have been chewed by the tobacco flea beetle, *Epitrix hirtipennis*.



Photo 9. Symptoms of tobacco flea beetle, *Epitrix hirtipennis*, on capsicum.



Photo 10. Symptoms of tobacco flea beetle, *Epitrix hirtipennis*, on potato.

Common Name

Tobacco flea beetle

Scientific Name

Epitrix hirtipennis

Distribution

Restricted. North, South (Guyana) and Central America, the Caribbean, Europe, Oceania. It is present in Fiji and Guam.

Hosts

Eggplant, and many others in the Solanaceae or nightshade family (capsicum, chilli, potato, tobacco, and tomato). Also, weeds, in the same family, e.g., *Physalis angulata*.

Symptoms & Life Cycle

Both the larvae and adults (Photos 1&2) damage plants: the adults feed on both sides of the leaves, chewing small, irregularly-shaped, holes in the leaves resulting in a 'shot hole' appearance (Photos 3-6&8-10). Usually, seedlings are most damaged. The larvae feed on roots.

Eggs are laid at the base of the stem of hosts. The larvae are slender, white with a brownish head, less than 5 mm long. They remain underground feeding on the small roots and root hairs.

Adults are hard-shelled, brown with black markings, about 1.5-2 mm long, with long, 12-segmented antennae. Wing covers have rows of fine but distinct punctures. The hind legs of adults are relatively large, and they jump like fleas.

Spread occurs on the wing, as flea beetles are strong fliers. Survival between crops occurs on weeds, and in the stalks of tobacco if they are left in the field after cutting off the leaves.

Impact

Large populations can kill or stunt seedlings, but economic damage of mature plants is rare for this species and other *Epitrix* flea beetles. Large numbers are more frequent (in Fiji at least) during periods of prolonged drought (Photo 7).

Although species of *Epitrix* flea beetles have been reported to feed on potato tubers, making tunnels in the flesh, in the US and Portugal, no damage of economic significance has been reported for *Epitrix hirtipennis*. However, analysis of the damage caused in Fiji, especially in times of drought may change this.

Detection & inspection

Look for the pits and tiny holes, many less than 1 mm, in leaves of seedlings and mature plants. Look for beetles, shiny brown and black in large numbers, especially during droughts.

Management

CULTURAL METHODS

Before planting:

- Remove weeds, especially those in the Solanaceae (potato or nightshade) family.
- Do not plant new crops next to those infested by the flea beetle.
- If seedlings are damaged in the field, use transplants produced in a nursery.
- Use mulches; these may interfere with activity of the stages in the soil.

During growth:

- Continue to remove weeds.
- Check for flea beetles twice weekly, especially after planting the transplants.
- Use protective covers over seedlings (e.g., floating row covers, or synthetic materials supported by hoops).

After harvest:

- Collect crop debris and burn or bury it.
- Do not plant consecutive susceptible crops; rotate with maize, brassicas or legumes of at least 2 years.

CHEMICAL CONTROL

If pesticides are necessary, use botanical (plant-derived pesticides) sprays first, as these may cause less harm to natural enemies, and cost less than synthetic commercial products.

- Use neem, derris, pyrethrum or chilli. If these are used, add soap to improve wetting the flea hoppers and covering them with sprays.
- Alternatively, use commercial biopesticides, e.g., spinosad (the product is called Success), or abamectin, derived from different bacteria. Spinosad has been tested against the eggplant flea beetle, *Epithix fuscula* in the US and found effective.
- On tobacco, several foliar insecticides have been recommended for the control of flea beetles. These have included synthetic pyrethroids (e.g. bifenthrin, cypermethrin, deltamethrin), carbamates, and neonicotinoids. On potato, in Europe, the neonicotinoid acetamiprid is used against a flea beetle damaging potato tubers (renewed until 2033).

AUTHOR Grahame Jackson & Mani Mua

Information from Peppers. Flea beetles. UC IPM. (<http://ipm.ucanr.edu/PMG/r604300611.html>); and Cuthbertson ACS (2015) Chemical and ecological control methods for *Epitrix* spp. *Global J. Environ. Sci. Manage* 1(1): 95-97 (http://www.gjesm.net/pdf_9553_82f0576dbd1b7d6b6eb5eed9b003d41.html); and from Flea beetles. Colorado State University Extension. (<http://extension.colostate.edu/topic-areas/insects/flea-beetles-5-592/>). Photo 1&2 Natasha Wright, Cook's Pest Control, Bugwood.org. Photos 3-6 Mani Mua, SPC, Sigatoka Research Station, Fiji. Photo 8 R.J. Reynolds, Tobacco Company Slide Set, R.J. Reynolds Tobacco Company, Bugwood2.org

Produced with support from the Australian Centre for International Agricultural Research under project HORT/2016/185: *Responding to emerging pest and disease threats to horticulture in the Pacific islands*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This fact sheet is a part of the app *Pacific Pests and Pathogens*

The mobile application is available from the Google Play Store and Apple iTunes.

