



# Pacific Pests and Pathogens - Mini Fact Sheet Edition

<https://apps.lucidcentral.org/ppp/>

## Sorghum midge (336)



Photo 1. Adult sorghum midge, *Stenodiplosis sorghicola*.




Diagram. Structure of a single sorghum spikelet.

### Summary

- Worldwide distribution. Sorghum (grain and forage), and wild grasses (especially Johnson grass). A major fly pest.
- Larvae eat the young seed heads.
- Midge, mosquito-like, orange, up to 2 mm long, clear wings, long antennae, and, for females, a long thin ovipositor. Eggs laid into flowers.
- Eggs hatch, larvae are white then orange, 2.5 mm long, spindle shaped. Pupae on the flower head. Some larvae go into resting phase, up to 5 years.
- Spread occurs when grain is moved with larvae in resting phase.
- Natural enemies: several parasitoid wasps, but only partially effective.
- Cultural control: plant early; weed grasses; increase seeding rate; rotate with sugar cane or peanuts; legume intercrops; burn stubble; resistant varieties.
- Chemical control: if >1-2 midges/head at flowering apply synthetic pyrethroids.

### Common Name

Sorghum midge

### Scientific Name

*Stenodiplosis sorghicola*; previously, *Contarinia sorghicola*.

AUTHOR Grahame Jackson

<sup>1</sup>Information from Swaine G (1971) *Agricultural Zoology in Fiji*. Her Majesty's Stationery Office. London; and CABI (2015) *Stenodiplosis sorghicola* (sorghum midge) Crop Protection Compendium ([www.cabi.org/cabi](http://www.cabi.org/cabi)); and from Sorghum midge. Department of Agriculture and Fisheries, Queensland Government. (<https://www.daf.qld.gov.au/plants/field-crops-and-pastures/broadacre-field-crops/integrated-pest-management/a-z-insect-pest-list/sorghum-midge>). Diagram Mississippi State University Extension. (<http://extension.msstate.edu/publications/information-sheets/sorghum-midge-control>). Photo 1 Alton N. Sparks, Jr., University of Georgia. Bugwood.org

Produced with support from the Australian Centre for International Agricultural Research under project PC/2010/090: *Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This mini fact sheet is a part of the app *Pacific Pests and Pathogens*

The mobile application is available from the Google Play Store and Apple iTunes.

