



## Pacific Pests, Pathogens & Weeds - Fact Sheets

<https://apps.lucidcentral.org/ppp/>

### Beet webworm (345)



Photo 1. Larva of beet webworm, *Spolodea recurvalis*.




Photo 2. Adult beet webworm, *Spolodea recurvalis*, showing white bands on wings and abdomen.



Photo 3. As in Photo 2, adult beet webworm, *Spolodea recurvalis*, showing white bands on wings and abdomen

#### Common Name

Beet webworm, Hawaiian beet webworm

#### Scientific Name

*Spolodea recurvalis*; previously known as *Hymenia recurvalis*.

#### Distribution

Worldwide. Asia, Africa, North, South and Central America, the Caribbean, Europe (restricted), Oceania. It is recorded from Australia, Cook Islands, Fiji<sup>1</sup>, New Caledonia, New Zealand, Samoa, Solomon Islands, Tonga, Tuvalu, Tonga, Wallis and Futuna.

#### Hosts

Mainly plants in the following families: Amaranthaceae (amaranthus), Fabaceae (beans, peanut, mung bean, soybean), Chenopodiaceae (beet, spinach), and Solanaceae (eggplant), and some weeds - amaranthus (pigweed) and chenopodium (lamb's quarters), and ornamentals (e.g., cockscomb).

#### Symptoms & Life Cycle

The larvae do the damage, and when numerous can defoliate crops rapidly. They also attack, flowers and pods. At first, it feeds only on the lower surface, leaving the uppersurface as clear 'windows'; later, it eats inside a tube formed by rolling leaves or drawing adjacent leaves together with the aid of silken threads.

Egg are laid singly or in groups near the midrib on the undersides of leaves, and hatch to produce a creamy-white larva with numerous

hairs (Photo 1). Mature larvae are up to 25 mm long, greyish-green, with a dark line down the middle of the back; they become reddish-pink before pupating in the soil inside a cocoon covered with soil particles. The adult is 10 mm long with a wingspan of 22-24 mm and characteristic white bands across the abdomen and wings (Photos 2&3). The life cycle is about 30 days.

Spread is by flight on the wing; the moth is famous for long distance migrations. Spread may also occur associated with the international trade in plants.

## Impact

The larva is especially important on Indian spinach (*amaranthus*) in Fiji, and in beet plantations in Australia. The damage can be serious in leguminous crops if it coincides with the time of pod filling. Economic damage is recorded on the ornamental cockscomb as the flowers can be easily damaged by just a few larvae.

## Detection & inspection

Look for the larvae rolled in silken cocoons in the folds of leaves: they have a distinctive dark line along the back. Look for the adults that shelter beneath leaves and take flight when disturbed; they are brown with white markings. The adults are attracted to light.

## Management

### NATURAL ENEMIES

There are many larval parasitoids, but there does not appear to have been programs to multiply and release them.

### CULTURAL CONTROL

Before planting:

- Do not plant next to crops that are infested with beet webworm.

During growth:

- Weed; especially remove wild amaranth and chenopodium species.
- Alternately, only remove the webbed leaves or press folded rolled leaves with finger and thumb to kill the larvae.

After harvest:

- Collect and destroy crop debris after harvest.

### CHEMICAL CONTROL

If pesticides are necessary, use botanical (plant-derived pesticides) sprays first, as these may cause less harm to natural enemies, and cost less than synthetic commercial products.

- Use neem, derris, pyrethrum or chilli. If these are used, add soap to help the chemical reach the caterpillars within the rolled leaves.
- Alternatively, use commercial biopesticides, e.g., spinosad (the product is called Success) or Bt - *Bacillus thuringiensis* var. *kurstaki*.
- If using Bt, note the following:
  - Ensure that Bt covers the plants: caterpillars will only die if they eat the Bt crystal proteins.
  - Eggs are not susceptible to Bt.
  - Bt should be used as soon as damage is seen.
  - Small larvae are more susceptible to Bt than fully grown ones.
- Synthetic pyrethroids are likely to be effective, but will also kill natural enemies, and should be avoided if possible.

---

AUTHOR Grahame Jackson

<sup>1</sup>Information from Swaine G (1971) *Agricultural Zoology in Fiji*. Her Majesty's Stationery Office, London; and CABI (2017) *Spoladea recurvalis* (Hawaiian beet webworm) Crop Protection Compendium. ([www.cabi.org/cpc](http://www.cabi.org/cpc)); and from Rice weevil. Department of Agriculture and Fisheries, Queensland Government. (<https://www.daf.qld.gov.au/plants/field-crops-and-pastures/broadacre-field-crops/integrated-pest-management/a-z-insect-pest-list/stored-grain-insect-pests/rice-weevil>). Photo 1 Alton N. Sparks, Jr., University of Georgia, Bugwood.org. Photo 2 Mark Dreiling, Bugwood.org. Photo 3 McCormack, Gerald (2007) Cook Islands Biodiversity Database, Version 2007.2. Cook Islands Natural Heritage Trust, Rarotonga. Online at: (<http://cookislands.bishopmuseum.org>).

This fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.

