



Pacific Pests, Pathogens & Weeds - Fact Sheets

<https://apps.lucidcentral.org/ppp/>

Chinese cabbage flea beetle (166)



Photo 1. Distinctive stripes of the Chinese cabbage flea beetle, *Phyllotreta undulata*.



Photo 2. Distinctive stripes of the Chinese cabbage flea beetle, *Phyllotreta undulata*.



Photo 3. Damage to Chinese cabbage by the Chinese cabbage flea beetle, *Phyllotreta undulata*.



Photo 4. Close-up of Chinese cabbage leaf showing holes and flea beetles, *Phyllotreta undulata*.

Common Name

Lesser striped flea beetle, small striped flea beetle, flea beetle, turnip flea beetle

Scientific Name

Phyllotreta undulata

Distribution

Worldwide. In the tropics, sub-tropics and temperate countries. In Oceania, it has been recorded from Australia, Fiji, New Caledonia and Vanuatu.

Hosts

Members of the cabbage family, Brassicaceae (Cruciferae), Chinese cabbage in particular.

Symptoms & Life Cycle

The adult beetles do the damage. They are 2 mm long, with a pair of wide yellow stripes on their wing cases (Photos 1&2). Adults feed on the leaves, and the larvae feed on the fine roots. Pupation takes place in small cocoons in the soil. The life cycle from egg to adult is about 3-4 weeks.

Spread of the beetle is by flight and wind over short distances, and by the domestic and international trade in fresh vegetables.

Impact

This is a very serious pest, especially on Chinese cabbage. The adult beetles feed on leaves by scraping, and by making many small holes (Photo 3&4). The Chinese cabbages become unfit for sale.

Growth of seedlings can be checked by the attack and even killed. Larger plants continue to grow and produce leaves, but when populations are high, the leaves are full of holes, and the plants become unmarketable.

The beetle can spread several virus diseases, e.g., *Turnip rosette virus*, *Turnip yellow mosaic virus* and *Turnip crinkle virus*. However, the virus is usually less of a problem than the direct feeding by the adult beetles.

Detection & inspection

Look for the pinprick-size scrapings and holes in the leaves. Look for tiny beetles (2 mm long) with two broad yellow stripes on their backs. They can be found within the whorl of new leaves and on the back of older ones.

Management

NATURAL ENEMIES

Nothing is known about the natural enemies of this pest in the South Pacific.

QUARANTINE

Countries that have not yet recorded the presence of *Phyllotreta undulata* will need to consider potential pathways for the introduction for this serious pest, and ensure that they do not allow its entry.

CULTURAL CONTROL

Before planting:

- Do not plant Chinese cabbages on the same land where they were harvested recently. *Phyllotreta* eggs and larvae will still be in the soil ready to emerge and infest the new crop.
- Do not plant crops near to those that are already infested with the beetles.
- Do not plant downwind from infested crops.

During growth:

- Use manure, compost or commercial fertilizers to accelerate seedling growth.
- Use sticky traps to monitor the presence of the flea beetle. These are rectangles of cardboard, coated with grease. Attach these to a wooden pole, e.g., a broom handle, and move it quickly along the rows of plants. As the flea beetles jump, they stick to the grease.

After harvest

- Collect and burn as much of the remains of the crop as possible when harvest is complete.
- Practice crop rotation. Replant with crops that are not in the cabbage family.

RESISTANT VARIETIES

Probably all varieties of Chinese cabbage are susceptible to attack by this beetle; however, head cabbages (green and purple varieties) are not attacked in Vanuatu.

CHEMICAL CONTROL

Such is the importance of this insect that chemical control may be necessary. Do the following:

- Use plant-derived products, such as derris, pyrethrum or chilli (with the addition of soap).
- Test local varieties of *Derris* (fish poisons) for their effectiveness as a spray. Note, *Derris* contains rotenone, so it should be used with caution.
- Alternatively, use synthetic pyrethroids as they are likely to be effective, but they will also kill natural enemies, if present.

This fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.

