



Pacific Pests, Pathogens & Weeds - Fact Sheets

<https://apps.lucidcentral.org/ppp/>

Citrus root & collar rot (264)



Photo 1. Root and collar rot of citrus caused by *Phytophthora nicotianae*.



Photo 2. Aboveground signs of root and collar rot of citrus caused by *Phytophthora nicotianae*.

Common Name

Citrus root and collar rot, *Phytophthora* foot and root rot (*Phytophthora nicotianae*), and brown rot of citrus fruit (*Phytophthora citrophthora*).

Scientific Name

Worldwide; two species are commonly found causing root and collar rots on citrus: *Phytophthora nicotianae*, and *Phytophthora citrophthora*.

Distribution

Worldwide. *Phytophthora nicotianae* is reported on citrus from Australia, Cook Islands, Federated States of Micronesia, Fiji, New Caledonia, Niue, Papua New Guinea, and Wallis & Futuna. It is also reported from Samoa, and Tonga on other hosts (**see Fact sheet nos. 149 & 154**). *Phytophthora citrophthora* is reported from Australia, and Fiji.

Hosts

Citrus root and collar rot (*Phytophthora nicotianae*) occurs on *bele* (**see Fact Sheet no. 149**), citrus, papaya (**see Fact Sheet no. 152**), passionfruit (**see Fact Sheet no. 154**), pineapple, tobacco, tomato (**see Fact Sheet no. 157**) and many other hosts. Brown rot of citrus fruit (*Phytophthora citrophthora*) occurs most commonly on citrus.

Symptoms & Life Cycle

Citrus root and collar rot diseases are common in wet areas (Photo 1). They cause a slow decline and death of citrus trees (Photo 2). The *Phytophthora* species involved are not fungi, although they have many fungal characteristics; they are water moulds or oomycetes, related to algae. They have motile spores and this sets them apart from fungi.

The two *Phytophthora* species cause similar diseases, and it is impossible to tell them apart on host symptoms. In general, *Phytophthora citrophthora* causes root and collar rots in cooler areas. There are also differences in appearance in culture, morphology and DNA tests.

The first sign of the diseases are spots on the trunk usually at the graft between the scion and rootstock or at soil level. The spots appear water-soaked, meaning they look wet, dark, usually sunken and greasy. The bark dies, dries and falls away, showing brown stained rots with cracks beneath. Often gum oozes from the wounds which can extend up to 50 cm above soil level and down to the roots. If the rot circles the trunk, the tree dies.

Root rots occur if soil moisture is high; in such situation the fibrous (fine) roots can be destroyed in a few days; the outer tissues become

soft and are easily stripped leaving only the white inner more resistant part, called the "stele". If the soil stays wet for a more than a few days, the larger roots can also be affected.

As root and collar rots progress, leaves turn yellow, dry and fall, and branches die back. Initially, symptoms may be more obvious on one side of the tree, corresponding to the part of the root system with the most damage.

Fruit may also become infected by spores splashing from the soil, developing a firm, leathery, brown rot with a strong smell of fermentation. Fallen fruit also become infected. Stored fruit may show a white cottony growth of the water mould, especially if kept at high humidity.

Infection depends on: (i) the bark remaining in contact with wet soil at ground level; (ii) soil and air temperatures of 26-32°C; (iii) wounded bark; (iv) susceptibility of the variety.

Survival of the water moulds occurs as thick-walled resting spores called 'chlamydospores'. When conditions are right (temperatures of 32-36°C, and chemical stimulants from the roots), the chlamydospores germinate and produce spores called 'sporangia'. Yet another spore is produced inside the sporangia called 'zoospores' (see **Fact Sheet no. 149**), and these are capable of swimming short distances before germinating and infecting. Zoospores are attracted to roots by the chemicals that they produce.

Spread of citrus water moulds occurs when sporangia, zoospores, or chlamydospores, reach stems and trunks at soil level providing water is in contact with bark for at least 5 hours. They are also splashed in rain drops from the soil to the lower fruit. Spread over longer distances is by the movement of water on the surface or within soil; it also occurs in soil on machinery and footwear, and also by the movement of contaminated nursery plants. The disease is worse in wet, heavy soils that do not drain rapidly after rain, or are prone to floods.

Impact

Citrus root and collar rots are serious diseases, and occur in all citrus growing areas in the humid tropics. The impact of the diseases depends on the soil type, the rainfall, and frequency of flooding, and the rootstocks used.

Detection & inspection

Look for dieback symptoms, often on one side of the tree, and inspect the trunk at soil level. Look for bark that is dry and dying or dead, and cankers (often sunken open wounds) exuding gum. A culture of the oomycete is required for identification; this can be obtained from the margin of trunk lesions or from the lesions on fruits.

Management

CULTURAL CONTROL

Cultural practices are important in the management of citrus root and collar rots.

Before planting:

- Avoid heavy, poorly drained soils, or dig drains or trenches to carry the water as quickly as possible away from the trees. Citrus requires good drainage!
- Bud high on root stocks (>30-45 cm) to avoid spores being splashed by rain onto susceptible parts of the stem or trunk.
- Ensure that nursery plants are free from *Phytophthora* species before field planting.

During growth:

- Keep weeds, including grass, away from the base of trees.
- Prune low hanging branches to at least 1 m above soil level.
- Frequently inspect trees, especially during the first 2 years after planting, and after flooding. Look for gum on the trunk. If infections are found, remove the affected bark and apply a fungal paste of, e.g., metalaxyl, phosphorus acid, or a copper fungicide.
- Remove soil (brought by ants) trapped behind tree guards on young trees, if these are being used.
- Avoid wounding the base of the trunks of the trees.
- Remove any trees with trunk infections that have led to their death.

RESISTANT VARIETIES

The choice of root stock varieties is very important in the management of citrus root and collar rot diseases. Which to use, however, depends on other diseases present, those caused by viruses and nematodes, especially. It also depends on the likelihood of the trees suffering from waterlogging, even if it only occurs occasionally.

Root stocks that have resistance to diseases include, trifoliate orange, sour orange, mandarin, and citrange hybrids. Swingle citrumelo, a hybrid between a grapefruit and a trifoliate orange is resistant to severe citrus tristeza, *Phytophthora* root rot, and has tolerance to waterlogging. Note that rough lemon and sweet orange are susceptible to *Phytophthora* root rots. It is important to ask local government experts for advice on the varieties to use.

In Fiji, the order of resistance to collar rot is trifoliate orange, sour orange, mandarin, sweet orange, grapefruit, bush lemon, lime, and Lisbon lemon.

CHEMICAL CONTROL

Apart from the use of fungicides to treat trunk cankers (see under Cultural Control above), their use is not recommended except in commercial production. Do the following:

- Apply foliar sprays of phosphorous acid, after times of flowering and the main leaf flushes. (Always refer to the product label for the correct method of application, timing, and also ways to avoid possible leaf burn.)
- Apply copper fungicides to protect lower fruit from spores in water splashed from the soil.
- Apply copper fungicides to the base of the trunk to prevent collar rots.

AUTHOR Grahame Jackson

Photo 1 Kohler F, Pellegrin F, Jackson G, McKenzie E (1997) *Diseases of cultivated crops in Pacific Island countries*. South Pacific Commission. Pirie Printers Pty Limited, Canberra, Australia. Photo 2 *Diseases of fruit crops in Australia* (2009). Editors, Tony Cooke, Denis Persley, Susan House. CSIRO Publishing. Information from Hardy S, Barkley P, Creek A, Donovan N (2012) Impacts and management of flooding and waterlogging in citrus orchards. NSW Government Department of Primary Industries; and from CABI (2015) *Phytophthora nicotianae* (black shank) and *Phytophthora citrophthora* (brown rot of citrus fruit) Crop Protection Compendium. (www.cabi.org/cpc).

Produced with support from the Australian Centre for International Agricultural Research under project PC/2010/090: *Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.

