



## Pacific Pests, Pathogens & Weeds - Fact Sheets

<https://apps.lucidcentral.org/ppp/>

### Rice bakanae disease (429)



Photo 1. Bakanae disease, *Gibberella fujikuroi*, as it appears in the field.




Photo 2. Comparison of symptoms: rice infected by bakanae disease, *Gibberella fujikuroi* (left), and healthy plant (right).



Photo 3. Roots from lower nodes on plant with bakanae disease, *Gibberella fujikuroi*.

#### Common Name

Bakanae disease of rice; also known as Fool's rice. Bakanae is from the Japanese for "foolish seedling". On maize it is known as maize Fusarium kernel rot (see **Fact Sheet no. 224**).

#### Scientific Name

*Gibberella fujikuroi*. The asexual state is *Fusarium moniliforme*. There are three mating groups.

#### Distribution

Widespread. Africa, Asia, North, South, and Central America, the Caribbean, Europe, Oceania. It is recorded from American Samoa, Australia, Fiji, Guam, New Caledonia, New Caledonia, Papua New Guinea, Solomon Islands, and Vanuatu.

#### Hosts

Rice, and many other crops, including: banana, cotton, cowpea, fig, maize, sorghum, sugarcane, and tomato. Although there are many hosts, those of most economic importance are in the Poaceae, the grass family. Grass weeds are also hosts.

#### Symptoms & Life Cycle

The disease is caused by a fungus that is seed and soilborne. Infected seed results in poor seedling emergence, and gaps in the rows. Surviving seedlings are yellow, thin, taller than usual with fewer tillers, and may wilt and die. Those that reach maturity become very tall

due to abnormal elongation of internodes, and do not form filled grain (Photos 1&2). On these plants, pink-white fungal growth occurs at soil level, and moves up the stems. The nodes on infected plants may be pink to purple beneath the leaf sheaths, and lower nodes may produce roots (Photo 3). Root rot occurs from soilborne infections.

Contamination of grain occurs at flowering by airborne spores – asexual ‘conidia’ and sexual ‘ascospores’ - formed on the tillers (stems and leaf sheaths). Infected seed develops into elongated, ‘bakanae’ plants when planted, and soil infestations lead to root rot. From the roots, the fungus travels up the tillers, producing spores, which spread to the new grain and to the soil. Spores are also produced in large numbers on the straw and stubble and contaminate the panicles at harvest.

Bakanae infection is favoured by dryland cultivation and high temperatures (>30°C). Lower temperatures and very dry soil favours root rot.

Spread occurs via spores carried on the wind or in water. Long-distance spread is on and in infected seed. The fungus remains viable in seed or about 1-2 years, and for about 1 year in the soil. Seedborne levels of up to 25% are recorded by IRRI.

## Impact

IRRI reports crop losses of up to 20% in outbreaks of bakanae in South and Southeast Asia [e.g., losses reported in Japan (20-50%), India (4%), and Thailand (15%)]. Graham states: "The disease is the most serious seed and soilborne disease of dryland rice in Fiji, particularly in the Rewa Valley of Viti Levu"<sup>1</sup>.

## Detection & inspection

Look for abnormally tall thin yellowish-green plants in the nursery and field. Look for white fungal growth at the base and lower parts of tillers. Strip away the leaf sheaths and look for pink to purple nodes on the stems. Look for roots at the nodes. Standard seed health protocols exist for this disease.

## Management

### CULTURAL CONTROL

Before planting:

- Rotate rice after 2 years with non-susceptible crops, such as vegetables or root crops.
- Use certified seed, free from contamination and infection. If unsure, use a seed dressing (see below).

During growth:

- Pull out infected plants as soon as they are seen.

After harvest:

- Burn straw and stubble, or plough into soil as soon as possible after harvest.

### CHEMICAL CONTROL

Treat seed with thiram or thiram combined with thiophanate-methyl to eradicate spores on the surface of seeds. Fungicides in the azole and strobilurin groups are also effective. Trials using bleach (sodium hypochlorite) have reduced the incidence of the disease.

*When using a pesticide, always wear protective clothing and follow the instructions on the product label, such as dosage, timing of application, and pre-harvest interval.*

---

AUTHOR Grahame Jackson

Information (and Photo 2) from Bakanae. Rice Knowledge Bank. IRRI. (<http://www.knowledgebank.irri.org/training/fact-sheets/pest-management/diseases/item/bakanae>); and CABI (2018) *Gibberella fujikuroi* (bakanae disease of rice) Crop Protection Compendium. (<http://www.cabi.org/cpc/>); Bakanae. Department of Primary Industries, NSW Government, Australia. (<https://www.dpi.nsw.gov.au/biosecurity/plant/insect-pests-and-plant-diseases/Bakanae>); and from Graham KM (1971) *Plant diseases of Fiji*. Her Majesty's Stationery Office. London. Photo 1&3 O.P. Sharma, NCIPM, New Delhi. India, Bugwood.org).

Produced with support from the Australian Centre for International Agricultural Research under project HORT/2016/185: *Responding to emerging pest and disease threats to horticulture in the Pacific islands*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

---

This fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.

