



Pacific Pests, Pathogens & Weeds - Fact Sheets

<https://apps.lucidcentral.org/ppp/>

Rice green leafhopper (417)



Photo 1. Adult rice green leafhopper, *Nephotettix nigropictus*.



Photo 2. Adult rice green leafhoppers, *Nephotettix* sp. (Nepal).



Photo 3. Rice plant with symptoms of rice tungro disease.

Common Name

Rice green leafhopper

Scientific Name

Nephotettix nigropictus; there are several other *Nephotettix* green leafhopper species that feed on rice, e.g., *Nephotettix malayanus* and *Nephotettix virescens*.

Distribution

Restricted. Asia, Africa (Cameroon), Oceania. It is recorded from Australia, Fiji, Guam, Federated States of Micronesia, Palau, and Papua New Guinea.

Hosts

Rice and wild grasses and sedges.

Symptoms & Life Cycle

The green leafhopper is more common in lowland rainfed and irrigated wetland rice than in upland rice. Nymphs and adults do the damage, directly by sucking the sap of rice plants, creating white patches on the tillers, and causing wilting and, if the population is high enough, they cause 'hopperburn' - a yellowing, browning and death of the plants. Indirectly, damage is caused by the green leafhopper

spreading virus diseases, e.g., rice tungro bacilliform virus, rice tungro spherical virus, and others. Note that tungro viruses are not present in Pacific island countries; nearest country is Irian Jaya.

Eggs are white, banana-shaped, laid in groups of about 15 into the leaf sheath or the midribs. The nymphs are yellow at first, turning light brown and then near black at maturity. The adults have two black spots, extending to the tips of the forewings, black tips to the wings, and black around the margins of the pronotum (extension from the head over the thorax) (Photo 1). The black bands on the head are less obvious in females.

Female leafhoppers disperse as the crop matures, but they are not strong fliers, traveling only about 1 km. Large migrations do not occur.

Impact

Rarely are green leafhoppers in sufficient numbers to cause severe damage directly; they feed more on grassy weeds than on rice; however, they transmit virus diseases to rice, even though they are considered to be less efficient in transmitting viruses than another green leafhopper, *Nephotettix virescens*.

Detection & inspection

Look for the black spots on the centres and tips of the wings, and, also, three parallel bands across the head, best seen from above. An entirely green form also exists, but black lines on the head distinguish it from other leafhoppers. Note, that regrowth of plants after harvest is a good time to determine the presence of disease.

Management

NATURAL ENEMIES

There are a large number of parasitoids and predators that attack all stages of the green leafhopper life cycle. As long as insecticides are not used, populations of this pest are usually under control by natural enemies. However, virus infection may still occur. Infection is often early in the crop, at times when parasitoids and predators are only beginning to increase, and numbers too small to control populations to the level required to prevent spread of viruses.

CULTURAL CONTROL

Before planting:

- Try to synchronise planting in any area so that crops do not overlap, to prevent pest populations moving from harvested to standing crops.
- Try to ensure that there is a rice-free period, and this coincides with the dry season (if there is one), so that hopper populations are low. During this time hoppers will lose the virus and their ability to infect. The ability of green leafhoppers to transmit tungro viruses is lost within 2 weeks. The virus is 'semi-persistent'.
- Avoid early infection by virus diseases: in irrigated systems. It is good practice to make nurseries away from rice fields and possible sources of infection.
- Plant when local knowledge suggests that crops will not coincide with seasonal peaks of leafhopper populations.
- Try planting older (>3 week-old) seedlings to delay the time that plants are exposed to infection in the vegetative stage.

During growth:

- Do not apply excessive amounts of nitrogen: follow local recommendations.
- Weed; remove grasses from around the fields, as these may be sources of virus.

After harvest:

- Monitor ratoon crops for virus disease. If present, establish new nurseries away from infected fields. Preferably, plough the land, turning the stubble into the soil.
- Rotate with other crops in the dry season: plant maize, soybean or peanut. Alternatively, fallow the land between rice crops.

RESISTANT VARIETIES

Most modern varieties released in Asia have resistance against *Nephotettix virescens*, and so they will be useful against *Nephotettix nigropictus*, too. Interestingly, resistant varieties force a change in feeding of the leafhopper from phloem to xylem, and as the virus is

concentrated in the phloem, fewer leafhoppers are infected.

CHEMICAL CONTROL

In Fiji, acephate and diazinon are recommended, but these insecticides should be used sparingly, and only when outbreaks of the leafhopper are likely to cause damage, which is relatively rare. Insecticides are more likely to result in the destruction of parasitoids and predators and cause increases of OTHER insects, e.g., the brown planthopper, and result in a rapid increase ('resurgence') of populations, leading to hopperburn.

Use neem as an alternative insecticide; it is effective and less harmful to natural enemies.

When using a pesticide, always wear protective clothing and follow the instructions on the product label, such as dosage, timing of application, and pre-harvest interval.

AUTHOR Grahame Jackson

Information from CABI Plantwise Knowledge Bank (<https://www.plantwise.org/KnowledgeBank/datasheet/36200>); and from Plant-sucking pests - green leafhoppers.

(<https://www.flickr.com/photos/ricephotos/6676383521/in/photostream/>). Photo 1 *Nephotettix nigropictus* Natasha wright, Cook's Pest Control, Bagwod.org. Photo 2&3 Bhusal K *et al.* (2019) A review of rice Tungro virus in Nepal. Journal of Plant Sciences and Crop Protection. Volume 2(1).

Produced with support from the Australian Centre for International Agricultural Research under project HORT/2016/185: *Responding to emerging pest and disease threats to horticulture in the Pacific islands*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.

