



Pacific Pests, Pathogens & Weeds - Fact Sheets

<https://apps.lucidcentral.org/ppp/>

Sugarcane red rot (221)



Photo 1. External red blotches on sugarcane stems caused by red rot, *Glomerella tucumanensis*.



Photo 2. Internal symptoms of red rot, *Glomerella tucumanensis*, in a sugarcane stem.



Photo 3. Red rot, *Glomerella tucumanensis*, showing symptoms of the fungus in the midrib of a sugarcane leaf. The small spots on the leaf blade are probably white rash. (see Fact Sheet no. 220.)

Common Name

Sugarcane red rot

Scientific Name

Glomerella tucumanensis; different strains are reported that vary in their ability to cause disease. The fungus is also known as *Colletotrichum fulcatum*; this produces spores without going through a sexual process.

Distribution

Worldwide. In the tropics. Asia, Africa, North, South and Central America, the Caribbean, Europe, Oceania. It is recorded from American Samoa, Australia, Cook Islands, Fiji, French Polynesia, Guam, Federated States of Micronesia, New Caledonia, Papua New Guinea, Samoa, Solomon Islands, Tonga, and Vanuatu.

Hosts

Sugarcane, other *Saccharum* species, maize and sorghum, and wild sorghum (*Sorghum halepense*).

Symptoms & Life Cycle

Stems show large red blotches on the outside (Photo 1), and internal red rots with white patches, when the stems are cut open (Photo 2). Cavities may be present which contain the cottony growth of the fungus. The leaves turn yellow, dry and die. As the rots develop, the stems are easily broken. Stems are infected through the nodes (where the leaf buds and roots develop) and through wounds.

On the leaves, small red oval spots occur on the midrib and upper surface of the leaf; these develop pale yellow to white centres, sometimes merging to cover the length of the leaf (Photo 3).

Spread of the fungus occurs in planting material, and this can be over both short and long distances. Spread also occurs as spores produced on infected plants, water-splashed to those nearby. The fungus can survive for a few months in the soil, on plant debris and in

sugarcane stems. Infection of the stems occurs through direct penetration of the nodes (buds, leaf scars or roots), from the fungus in the soil or in rain/irrigation water.

Cool, wet weather, heavy soils, and continuous planting of the same variety in the same land favour the disease. Drought also increases susceptibility to the disease.

Impact

The fungus attacks setts in the soil, causing them to rot as they sprout, and it also infects the standing crop causing plants to wilt and shrivel. In addition to the loss of plants, there is a reduction of sucrose content in those that are affected by rots. Severe losses in yield of cane and, consequently, sucrose have been reported from Bangladesh, India, Pakistan and Thailand, and it is likely that similar losses occur elsewhere.

Detection & inspection

Look for red blotches on the stems and elongated spots in the midribs on the top surface of leaves. Look for white spots or blotches in the pith which are usually elongated at right angles to the long axis of the stem; these are diagnostic for this disease.

Management

CULTURAL CONTROL

Before planting:

- Use healthy setts; take planting material from fields where the disease has not been seen.
- Plant when either wet or dry times will be avoided.
- If growers have only a small number of setts, treat with hot water at 50°C for 2 hours; obviously, this is not an appropriate method for large amounts of planting material.
- Avoid heavy soils which have a tendency to become waterlogged. Red rot is reported to be worse where poor drainage and infection by *Pythium* (an oomycete) occur together.

After harvest:

- Collect the remains of the crop and burn it.
- Do not ratoon diseased crops, otherwise the fungus may increase to an epidemic.
- Plough the field several times to expose the fungus to sunlight.
- Do not plant one crop of sugarcane after another in the same land; use a rotation of at least 2-years.

RESISTANT VARIETIES

There are varieties resistant to this disease; check those available in your area. There are, however, a number of strains of this fungus.

CHEMICAL CONTROL

This is not a method that can be used to control this disease economically.

AUTHOR Grahame Jackson

Information (and Photos 1&2) CAB International (2014) *Glomerella tucumanensis* (red rot of sugarcane) Crop Protection Compendium. (<http://www.cabi.org/cpc/>). Photo 3 Kohler F, Pellegrin F, Jackson G, McKenzie E (1997) *Diseases of cultivated crops in Pacific Island countries*. South Pacific Commission. Pirie Printers Pty Limited, Canberra, Australia.

Produced with support from the Australian Centre for International Agricultural Research under project PC/2010/090: *Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.





Australian Government
Australian Centre for
International Agricultural Research

Copyright © 2020. All rights reserved.