



## Pacific Pests, Pathogens & Weeds - Fact Sheets

<https://apps.lucidcentral.org/ppp/>

### Sweetpotato scurf (260)



Photo 1. Dark grey patches on the surface of sweetpotato storage roots caused by scurf, *Monilochaetes infuscans*.



Photo 2. Scurf, *Monilochaetes infuscans*, symptoms on variety Beauregard.

#### Common Name

Sweetpotato scurf

#### Scientific Name

*Monilochaetes infuscans*

#### Distribution

Worldwide. Asia, Africa, North (Hawaii), South America, Europe, Oceania. It is not recorded in Fiji, Samoa, and Solomon Islands. It is recorded from Cook Islands, Federated States of Micronesia, New Zealand, Northern Mariana Islands, Palau, Papua New Guinea, Tonga, and Vanuatu.

#### Hosts

Sweetpotato, and wild *Ipomoea* species.

#### Symptoms & Life Cycle

A fungus causes the disease. Greyish spots and patches (these may be purplish-brown to black on orange fleshed varieties) occur, with irregular margins, joining together and often covering most of the storage roots (Photos 1&2). The colour of the infections depends on the variety. However, the infections are only in the "periderm" of the storage roots; this is the thin "skin" outside the starchy flesh. Spores are produced on the surface of the infected spots and patches.

Spots, similar to those on the storage roots, may occur on the stems and leaves if infected storage roots are used as a source of planting material.

The fungus is worse in soils with high organic matter, and much less in light sandy soils. Also, scurf is worse during rainy seasons. Spread occurs on infected cuttings, and these can be a source of spores to infect the storage roots. Survival occurs in crop debris for 2-3 years.

#### Impact

Scurf is a 'cosmetic' disease, meaning it does not reduce the yield of the storage roots or their taste, but it affects their appearance, which could mean a lower price in the market.

#### Detection & inspection

Look for greyish spots and patches on the storage roots; they are especially noticeable on white skinned varieties. They are darker on orange-fleshed varieties (Photo 2).

## Management

### CULTURAL CONTROL

The following will help to reduce the amount of infection:

Before planting:

- Use a crop rotation of 2-3 years for sandy loam soils - longer (3-4 years) for soils high in organic matter.
- Remove wild *Ipomoea* weeds, i.e. those belonging to the morning glory family.
- Make sure cuttings for planting are free from infection: (i) take cuttings only from plants whose storage roots were free from symptoms at harvest; (ii) if using storage roots to grow sprouts for planting, choose only those free from symptoms of scurf. It is easier to see symptoms if the storage roots are wet. Cut sprouts (also called "slips") 2-3 cm above soil level to avoid carrying the fungus to the field.
- Use land free from the scurf fungus, i.e. land where previous harvests of storage roots have been free from symptoms.
- Do not add organic manures to soil where scurf symptoms occur on storage roots.

After harvest:

- Remove harvested vines and burn them. Do not incorporate into the soil.

### RESISTANT VARIETIES

There are no varieties that are resistant to the fungus. Varieties that form storage roots lower in the soil may have less infection because they are further from the source of spores.

### CHEMICAL CONTROL

There are no fungicides that can be used to manage this disease.

---

AUTHOR Grahame Jackson

Information from CABI (2015) Crop Protection Compendium *Monilochaetes infuscans* (scurf of sweet potato). ([www.cabi.org/cpc](http://www.cabi.org/cpc)).

Produced with support from the Australian Centre for International Agricultural Research under project PC2010/090: *Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

---

This fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.

