



Pacific Pests, Pathogens & Weeds - Fact Sheets

<https://apps.lucidcentral.org/ppp/>

Taro hornworm (032)



Photo 1. A caterpillar of taro hornworm, *Hippotion celerio*, eating a leaf and leaving only the veins.



Photo 2. Severe damage to taro caused by the taro hornworm, *Hippotion celerio*; the leaves have been eaten leaving only the leaf stalks or petioles.




Photo 3. Solitary egg of taro hornworm, *Hippotion celerio*.



Photo 4. When mature the caterpillars of taro hornworm, *Hippotion celerio*, are dark brown.



Photo 5. Pupa taro hornworm, *Hippotion celerio*.



Photo 6. Adult taro hornworm, *Hippotion celerio*.

Common Name

Taro hornworm

Scientific Name

Hippotion celerio

Distribution

Widespread. In tropical, sub-tropical and temperate countries. Asia, Africa, Europe, Oceania. It is recorded from Australia, Fiji, Guam, Kiribati, Nauru, New Caledonia, New Zealand, Papua New Guinea, Samoa, Solomon Islands, Tonga, and Vanuatu.

Hosts

Taro, sweetpotato, and noni (*Morinda* sp.)

Symptoms & Life Cycle

The caterpillars do the damage. They eat the leaves, leaving only the main veins (Photo 1). Sometimes even these are eaten and only the

petioles are left (Photo 2).

The eggs, round, 1-1.5 mm diameter, are laid singly on both sides of the leaf and on the petioles (Photo 3). They vary from round to oval, and from clear to bluish-green. Just before the emergence of the caterpillars they are green-yellow. The caterpillars moult four times: at first they are pale yellow, then, as they age, they become green and dark brown (Photo 4). Some remain green. The eyespots and a yellow line running almost the entire length of the caterpillar develop after the first moult. When mature, the caterpillars are 80-90 mm long. At this stage, they move to the soil, form a cocoon or cell in the leaf litter or just below the soil, and pupate (Photo 5). The pupae are grey-brown, 45-50 mm long, with dark brown specks. This period lasts 15-18 days.

Adults have wingspans of 40-90 mm; they are streamlined, with large heads and eyes (Photo 6). The forewings are long, narrow and much larger than the hind wings. The forewings have an oblique silver stripe across a generally grey background, while the hind wings may have pink or brown and black areas. At rest, the wings are 'tented' over the body. The moths are capable of flying long distances and are attracted to light.

Impact

Outbreaks are uncommon as the caterpillars are under control from natural enemies; otherwise, the caterpillars can be very damaging.

Detection & inspection

The damage done to taro is characteristic of this hornworm. Look for the way that the leaves are eaten between the veins. Also, the caterpillars can be found on the leaves during the day, especially on the underside. Look for a large caterpillar with eyespots at the front end, with a line along the sides, and a horn at the end.

Management

NATURAL ENEMIES

These include egg parasites - *Trichogramma* sp. (a wasp); and larval parasites *Palexorista* sp. (a fly), *Snellenius hippotionus* (a wasp), and there are probably many more. The cane toad may also feed on the larvae.

CULTURAL CONTROL

It is important that growers visit the gardens frequently and regularly, twice a week is recommended, to check if there are egg masses and young caterpillars on the leaves. Mostly, taro hornworm is under control by natural enemies, and it is only very occasionally that outbreaks occur.

The caterpillars are large, and handpicking is an option if the numbers and areas of infestation are small. If they are large, chickens are often a good way of managing outbreaks.

CHEMICAL CONTROL

Pesticides are not normally recommended for the control of this moth on taro. They are only needed when the natural enemies have been destroyed by cyclones, droughts, or when plantings are in isolated places. In these situations, do the following:

- Use plant-derived products, such as neem, derris, pyrethrum and chilli (with the addition of soap), or commercial products that contain disease-causing organisms, such as spinosad (Success) and Bt - *Bacillus thuringiensis* subspecies *kurstaki*.
- Note, a variety of *Derris*, brought many years ago to Solomon Islands from Papua New Guinea, is effective as a spray. It contains rotenone, an insecticide, so it should be used with caution. There may be varieties of *Derris* (fish poisons) in your country that can be tried (see **Fact Sheet no. 56**).
- Synthetic pyrethroids are likely to be effective, but will also kill natural enemies.

AUTHORS Helen Tsatsia & Graeme Jackson

Photo 3 Mike Furlong, University of Queensland, Brisbane. Photo 5 Wikipedia. (http://en.wikipedia.org/wiki/Hippotion_celerio). Photo 5 Mani Mui, SPC, Sigatoka Research Station, Fiji.

Produced with support from the Australian Centre for International Agricultural Research under project PC/2010/090: *Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.

