Pacific Pests, Pathogens & Weeds - Mini Fact Sheet Edition

https://apps.lucidcentral.org/ppp/

Breadfruit (fluted scale) mealybug (184)



Photo 1. Adult breadfruit mealybugs, *Icerya aegyptiaca*, showing the long white waxy fringe around the body.



Photo 2. Underside of adult breadfruit mealybug, *Icerya aegyptiaca*, to show the legs and antennae.

Summary

- Widespread distribution. Asia, Africa, Oceania. Principally breadfruit but also avocado, banana, citrus, jackfruit, mango, soursop, taro, and many others, including ornamentals. An important pest.
- Eggs laid into a sac producing "crawlers" that spread the pest. After a day, they settle down, feed, moult, becoming adult with 3-4 mm waxy processes.
- Damage done: (i) sucking sap, (ii) leaves turn black fungi grow on honeydew excreted by mealybugs.
- Natural enemies: ladybird beetles (Rodolia) important
- Cultural control: prune infested stems, branches and fruits, and burn; add fertilizer, manure or compost to improve tree health; assist natural enemies: destroy ants nests (hot water), prune low branches, and weed.
- Chemical control: soap, white or horticultural oils; PDPs: neem, derris, pyrethrum, or chilli; if ants are present use synthetic pyrethroids. Priority is to remove ants to let natural enemies work.

Common Name

Breadfruit mealybug, Egyptian fluted scale, Egyptian mealybug

Scientific Name

Icerya aegyptiaca

AUTHOR Grahame Jackson

¹Waterhouse DF (1993) Biological Control Pacific Prospects - Supplement 2. ACIAR Monograph No. 20. Brown Prior Anderson, Barwood, Victoria; and from CABI (2015) Icerya aegyptiaca. Crop Protection Compendium. (http://www.cabi.org.cpc/). Photo 1 Alessandra Rung, California Department of Food & Agriculture, Bagwood.org. Photo 2 Peter Ooi, Department of Agriculture & Food Sciences, Faculty of Science, Universiti Tunku Abul Rahman, Jalan University, Malaysia.

Produced with support from the Australian Centre for International Agricultural Research under project PC/2010/090: Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This mini fact sheet is a part of the app Pacific Pests, Pathogens & Weeds

The mobile application is available from the Google Play Store and Apple iTunes.



Copyright © 2020. All rights reserved.