

Pacific Pests, Pathogens & Weeds - Mini Fact Sheet Edition

https://apps.lucidcentral.org/ppp/

Cabbage grey leaf spot (310)



Photo 1. Cabbage grey leaf spot, Alternaria brassicae



Photo 2. Grey leaf spot, *Alternaria brassicae*, on cabbage. Note the concentric rings typical of *Alternaria* infections.



Photo 3. Grey leaf spot, *Alternaria brassicae* on cabbage.




Photo 4. Spores of grey leaf spot, *Alternaria brassicae*.

Compare with spores of *Alternaria brassicicola* (see

Fact Sheet 319).

Summary

- Worldwide distribution. On plants in the cabbage family, especially Chinese cabbage and mustards.
 Important fungal disease, simlar to cababge black leaf spot (see Fact Sheet no. 133).
- Damage: i) seed infections cause pre- and post-emergence damping-off (see Fact Sheet no. 47); ii) seed infection; and iii) spots on head and/or outer leaves unsightly reducing quality and shelf life.
- Grey leaf and stem spots, circular or irregular, mostly between veins, up to 25 mm diameter, with concentric rings, producing spores below. Centre of spots fall out with age.
- Dark, sunken rots occur on heads of broccoli and cauliflower.
- Spread by water splash, in wind-driven rain, on seed.
- Cultural control: seed hot water (50°C for 25-30 mins.); nursery hygiene: clean trays, pasteurise soil or use soilless mixes, and
 discard infected seedlings; weed; space plants to allow air movement; avoid overhead irrigation; collect and destroy debris after
 harvest; crop rotation.
- Chemical control: seed captan or thiram; field chlorothalonil, copper or mancozeb.

Common Name

Cabbage grey leaf spot

Scientific Name

Alternaria brassicae. Another Alternaria fungus, Alternaria brassicicola, cabbage black leaf spot, also occurs, and causes similar

symptoms (see Fact Sheet no. 133). Microscopic examination of the spores is needed to distinguish between the two species (Photo 4).

AUTHOR Grahame Jackson and Eric McKenzie

Information from Diseases of vegetable crops in Australia (2010). Editors, Denis Persley, Tony Cooke, Stsan House. CSIRO Publishing and from Gerlach WWP (1988) Plant diseases of Western Samoa. Samoan German Crop Protection Project, Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ) Gribh, Germany. Photo 1 Gerald Holmes, California Polytechnic State University at San Luis Obispo, Bugwood org. Photo 2 Yuan-Min Shen, Taichung District Agricultural Research and Extension Sation, Bugwood org. Photo 3 Howard F. Schwartz, Colorado State University, Bugwood org. Photo 4 (taken by Eric McKenzie), and used in this fact sheet, appeared previously in McKenzie E (2013) Alternaria brassicae PaDIL - (http://www.padil.gov.au).

Produced with support from the Australian Centre for International Agricultural Research under project PC2010/090: Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This mini fact sheet is a part of the app Pacific Pests, Pathogens & Weeds

The mobile application is available from the Google Play Store and Apple iTunes.









Copyright © 2020. All rights reserved.