



Pacific Pests, Pathogens & Weeds - Mini Fact Sheet Edition

<https://apps.lucidcentral.org/ppp/>

Elephant's foot (459)

Relates to: **Weeds**




Photo 1. Stand or thicket of elephant's foot, *Elephantopus mollis*.



Photo 2. Rosette of leaves, elephant's foot, *Elephantopus mollis*.



Photo 3. Flowerhead, elephant's foot, *Elephantopus mollis*.



Photo 4. Close-up flowerhead, elephant's foot, *Elephantopus mollis*. Note the leaf-like bracts under the flowers.

Summary

- Widespread. Asia, Africa, North, South and Central America, Caribbean, Oceania. In many Pacific islands.
- Aggressive perennial weed of plantations, forming dense stands in sunny, dry, disturbed places, in pastures, where rainfall moderate to high, and threat to native species. Seeds prolifically and leaves of young plants produce thick ground cover smothering competitive species. Cattle do not eat it.
- Fast-growing, up to 1.5 m tall. Forms rosette of leaves, each up to 30 cm long, underside sticky and covered with hairs. One or more branched, flowering stems with smaller, alternate leaves, toothed margins, and winged leaf stalks. Flowerheads white flowerheads up to 2 cm wide, in clusters. Seeds black with bristles.
- Spread: seeds, by fur of animals, clothing, wind; contaminant of pasture seeds; perhaps for medicine and as ornamental.
- Biosecurity: Among 10 most important weeds of Vanuatu. On Global Invasive Species Database of alien invasive species (IUCN, 2020). Noxious weed in Fiji.
- Biocontrol: Fly, *Tetreauresta obscuriventris*, introduced to Fiji and Hawaii. Little impact.
- Cultural control: Hoeing; mowing (plus herbicide); plough, disc and harrow; vehicle hygiene.
- Chemical control: in Australia: fluroxypy; glyphosate; triclopyr + picloram; metsulfuron-methy; picloram. Elsewhere, glyphosate (e.g., Fiji); MCPA (e.g., Fiji); 2,4-D.

Common Name

Elephant's foot; it is also known as tobacco weed.

Scientific Name

Elephantopus mollis. Some authorities consider it synonymous with *Elephantopus scaber*, but others keep them separate, although they are closely related. It is a member of the Asteraceae.

AUTHORS Grahame Jackson & Aradhana Deesh

Adapted from Elephant's foot (*Elephantopus mollis*) (2018) Weeds of SE Qld and Northern NSW. Lucidcentral. (<https://www.lucidcentral.org/editors-pick-animal-and-plant-identification-keys/key-to-weeds-of-se-qld-and-northern-nsw/>); and additional information from CABI (2019) *Elephantopus mollis* (elephant's foot). Invasive Species Compendium. (<https://www.cabi.org/isc/datasheet/41496#tosummaryOfInvasiveness>); and Waterhouse DF, Norris KR (1987) *Biological Control Pacific Prospects*. Inkata Press, Melbourne. Photo 1 Forest & Kim Starr. Starr Environmental, bugwood.org. Photo 2 Vinayaraj Elephantopus_scaber. Photo 3 Gerald McCormack, Cook Islands Biodiversity & Natural Heritage. (<http://cookislands.bishopmuseum.org>). Photo 4 OM Montiel Close-up of the flowerhead.

Produced with support from the Australian Centre for International Agricultural Research under project HORT/2016/185: *Responding to emerging pest and disease threats to horticulture in the Pacific islands*, implemented by the University of Queensland, in association with the Pacific Community and Koronivia Research Station, Ministry of Agriculture, Fiji.

This mini fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.

