

Pacific Pests, Pathogens & Weeds - Mini Fact Sheet Edition

https://apps.lucidcentral.org/ppp/

Nutgrass (453)

Relates to: Weeds



Photo 1. Mature plant, nutgrass, *Cyperus rotundus*, with adjacent 'daughter' plants arising from underground tubers.



Photo 2. Single mature plant, nutgrass, *Cyperus* rotundus, to show underground stem or rhizome, with



Photo 3. Triangular flower stem, nutgrass, *Cyperus rotundus*.



Photo 4. Flowerhead, nutgrass, *Cyperus rotundus*, with branches, each with flattened reddish-brown spikelets, which contain the many flowers. Note, the leaf-like

Summary

- Widespread. Asia, Africa, North, South and Central America, Caribbean, Europe, Oceania. In most Pacific countries.
- Invasive, grows very rapidly in a wide variety of environments, irrespective of soil types. Problematic in field crops, orchards, vineyards, pastures, gardens, fallows, natural and waste areas, footpaths, roadsides, banks of waterways, produces underground tubers that regenerate plants, and is tolerant of many herbicides.
- Plants 20-60 cm tall. Leaves, narrow, up to 20 cm long, at base of plants. Stems, underground, producing small tubers in chains, white and fleshy at first, later brown and wiry. Buds on the tubers form new plants. Flower stems, smooth, glossy, triangular in cross-section, with 3-8 branches of different lengths and leaf-like structures beneath; each branch bears several reddish-brown 'spikelets' with flowers. The fruits are 3-angled, brown or black, with a single seed inside. Germination very low, but many seed produced.
- Spread: underground stems; by cultivation and machinery; seed (probability low); flood waters; as a specialist food.
- Biosecurity: high risk of introduction; one of the worst 10 weeds in Fiji, French Polynesia, Guam, New Caledonia, PNG, Solomon
 Islands, Tonga. On Global Invasive Species Database of alien invasive species (IUCN, 2020). Biocontrol: a moth, *Bactra* species,

and a mycoherbicide, using Dactylaria higginsii.

- Cultural control: hand weed; shade; plough and disc; polyethylene mulch; vehicle hygiene to remove tubers and seed; ducks and/or chickens will eat the tubers.
- Chemical control: in Australia: glyphosate; 2,4-D; and the selective herbicide, halosulfuron-methyl. In Fiji, MCPA.

Common Name

Nutgrass; CABI prefers the name purple nutsedge.

Scientific Name

Cyperus rotundus. It is a member of the Cyperaceae.

AUTHORS Grahame Jackson & Aradhana Deesh

Adapted from Nutgrass (Cyperus rotundus) (2018) Weeds of SE Qld and Northern NSW. Lucidcentral. (https://www.lucidcentral.org/editors-pick-animal-and-plant-identification-keys/key-to-weeds-of-se-qld-and-northern-nsw); and additional information from CABI (2019) Cyperus rotundus (purple nutsedge). Invasive Species Compendium. (https://www.cabi.org/ise/datasheet/17506); and Waterhouse DF, Norris KR (1987) Biological Control Pacific Prospects. Inkata Press, Melbourne; and from Coleman Met al., (2018) Nutgrass (Cyperus rotundus): Weed management guide for Australian vegetable production. School of Environmental and Rural Science, University of New England, Armidale. (https://www.une.edu.au/_data/ssets/pdf_file/0006/235995/une-weeds-nutgrass.pdf). Photos 1&3 Rickipelleg Nutgrass - Cyperus rotundus. Photo2 Joseph LaForest, University of Georgia, Bugwood org Photo 4 Cyperus rotundus by_Jevan Jose, kadavoor, Kerala, India.

Produced with support from the Australian Centre for International Agricultural Research under project HORT/2016/185: Responding to emerging pest and disease threats to horticulture in the Pacific islands, implemented by the University of Queensland, in association with the Pacific Community and Koronivia Research Station, Ministry of Agriculture, Fiji.

This mini fact sheet is a part of the app Pacific Pests, Pathogens & Weeds

The mobile application is available from the Google Play Store and Apple iTunes.









Copyright © 2020. All rights reserved.