



Red imported fire ant (363)

Relates to: [Ants](#)



Photo 1. Front view, 'minor' worker, red imported fire ant, *Solenopsis invicta*.



Photo 2. Side view, 'minor' worker, red imported fire ant, *Solenopsis invicta*.



Photo 3. Gaster, 'minor' worker, red imported fire ant, *Solenopsis invicta*.



Photo 4. Front view, 'major' worker (soldier), red imported fire ant, *Solenopsis invicta*.



Photo 5. Side view, 'major' worker (soldier), red imported fire ant, *Solenopsis invicta*.



Photo 6. Gaster, 'major' worker (soldier), red imported fire ant, *Solenopsis invicta*.

Summary

- Worldwide distribution. Not recorded from Pacific island countries. Present in Australia (under eradication). Among world's 100 worst invasive species.
- Ground-nesting in sunny agricultural, forestry, grasslands, urban areas. More aggressive and painful sting than tropical fire ant.
- Nests - mounds up to 40 cm, 30-50 cm diameter; entrance holes 5-10 m away. Eat seeds, other insects, smaller vertebrates and honeydew.
- Direct damage: vegetable seeds, fruits, new shoots and roots (large annual costs in US and Australia); loss of biodiversity; indirectly by defending aphids, mealybugs, scales and whiteflies from natural enemies.
- Queens, males and three types of workers: minor, major and intermediate. Nests mostly with 2 or more queens, Up to 500 colonies/ha and each colony 200-400,000. Some with single queens.
- Tramp ant. Spread by queens leaving nest, nuptial flight then new site, or 'budding'; spread via horticultural trade.

- Biosecurity: requires risk assessments, regulations preventing introduction, protocols in case of breaches, and ability to make rapid response. Pacific Ant Prevention Plan available (IUCN/SSC Invasive Specialist Group).
- Natural enemies: several *Pseudacteon* species under study in US.
- Cultural control: hot water at 47°C kills ants; over 49°C kills plants.
- Chemical control: use i) stomach poisons (fibronil, Amdro borax), ii) growth regulators (methoprene, pyriproxyfen), iii) nerve poisons (bifenthrin, fipronil, imidacloprid). See (<http://piat.org.nz/getting-rid-of-ants>).

Common Name

Red imported fire ant

Scientific Name

Solenopsis invicta

AUTHOR Grahame Jackson

Information from CABI *Solenopsis invicta* (red imported fire ant) (2017) Crop Protection Compendium (www.cabi.org/cpci); and Pacific Invasive Ant Toolkit. (<http://www.piat.org.nz/problem-ants/worst-5-identification/red-imported-fire-ant>); and *Solenopsis invicta*. AntWiki. (http://www.antwiki.org/wiki/Solenopsis_invicta); and AntWeb. (<https://www.antweb.org/description.do?genus=solenopsis&species=invicta&rank=species&resetProject=true>); and from *Solenopsis invicta* (2018) Global Invasive Species Database. (<http://www.iucngisd.org/gisd/species.php?sc=77>). Photos 1-6 Amy Carmichael (2005) Red imported fire ant (*Solenopsis invicta*). Queensland University of Technology. PaDIL - (<http://www.padil.gov.au>).

Produced with support from the Australian Centre for International Agricultural Research under project HORT/2016/185: *Responding to emerging pest and disease threats to horticulture in the Pacific islands*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This mini fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.

