



Pacific Pests, Pathogens & Weeds - Mini Fact Sheet Edition

<https://apps.lucidcentral.org/ppp/>

Rice white stem borer (411)



Photo 1. Adult rice white stem borer, *Scirpophaga innotata*.



Photo 2. Adult rice white stem borer, *Scirpophaga innotata*.



Photo 3. Damage ('deadheart') to rice stem by *Chilo auricilius* (damage to *Scirpophaga innotata* is similar).



Photo 4. 'Whitehead' - a symptom caused by stem borers: the base of the panicle is damaged preventing it from emerging or, if already emerged, the grain is unfilled and white.

Summary

- Restricted. Southeast Asia, Oceania. In Australia, Papua New Guinea.
- Severe on rice and wild rice if attack comes at flowering stage.
- Larvae tunnel between stem and leaf sheaths to the growing point, killing it; stems pull out easily ('deadhearts'). Panicles fail to emergence, or emerge with white unfilled grain ('whiteheads').
- Eggs laid up to 100 near leaf tips, covered in hairs. Larvae white to yellowish when mature, 25 mm long. Pupae white. Adults white, wingspans 18-33 mm (males smaller than females). Nocturnal.
- Note, larvae go through resting period (diapause) for several months if conditions unfavourable. Where crops one a year and long-maturing, diapause means many moths emerge at one time.
- Natural enemies: many egg and larval parasitoids and predators.
- Biosecurity: introduction possible on produce contaminated with infested stems of host plants.
- Cultural control: plough land well (IMPORTANT to bury larvae/pupae of previous crop) and direct seed; cut tips of seedling to remove eggs; plant at higher density than normal; rotate, e.g., legumes; synchronise plantings with neighbours; submerge eggs by raising water occasionally; weed; apply split applications N; harvest at ground level to remove larvae; plough in stubble, unharvested plants and weeds; in irrigated rice-rice-fallow systems use medium (135-140 day) varieties so diapause is incomplete when time to replant.
- Chemical control: unlikely to be needed. Use abamectin. Avoid broad-spectrum insecticides to preserve natural enemies.

Common Name

Rice white stem borer

Scientific Name

Scirpophaga innotata. A moth in the Crambidae.

AUTHOR Grahame Jackson

Information (and Photo 4) from Rice Knowledge Bank, IRRI (<http://www.knowledgebank.irri.org/training/fact-sheets/pest-management/insects/item/stem-borer>); and CABI *Scirpophaga innotata* (white rice stem borer) (2017) Crop Protection Compendium. (www.cabi.org/cpc); and from Pathak MD, Khan ZR (1994) Insect Pests of Rice. IRRI/ICIPE. Photo 1 Biodiversity India. Rice white stem-borer moth. (<https://indiabiodiversity.org/observation/show370837>). Photo 2 11ANIC-04905 CSIRO/BIO Photography Group, Centre for Biodiversity Genomics. (http://v3.boldsystems.org/index.php/Taxbrowser_Taxonpage?taxid=211612). Photo 3 Anderson S, Tran-Nguyen L (2012) Gold-fringed Rice Borer (*Chilo auricilius*). (Source: N. Sallam DAFF Biosecurity.) PaDIL - (<http://www.padil.gov.au>);

Produced with support from the Australian Centre for International Agricultural Research under project HORT/2016/185: *Responding to emerging pest and disease threats to horticulture in the Pacific islands*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

This mini fact sheet is a part of the app *Pacific Pests, Pathogens & Weeds*

The mobile application is available from the Google Play Store and Apple iTunes.

