

# Pacific Pests, Pathogens & Weeds - Mini Fact Sheet Edition

https://apps.lucidcentral.org/ppp/

## Sensitive plant (464)

Relates to: Weeds



Photo 1. Prostrate (lying flat) habit, sensitive plant, *Mimosa pudica*.



Photo 2. Thorns on the stem of sensitive plant, *Mimosa pudica*.



Photo 3. Leaves and leaflets, sensitive plant, *Mimosa* pudica



Photo 4. Leaves and leaflets, sensitive plant,  $\it Mimosa$   $\it pudica$ .



Photo 5. Sensitive plant, *Mimosa pudica*. On some of the leaflets, the bristles at the edges can just be seen.



Photo 6. Flowerheads, sensitive plant, Mimosa pudica.



Photo 7. Flowerhead, sensitive plant, Mimosa pudica.



Photo 8. Fruits divided into 1-seeded segments sensitive plant, *Mimosa pudica*.

## **Summary**

- Widespread. Asia, Africa, North, South and Central America, Caribbean, Oceania. In many Pacific islands.
- Thrives in disturbed areas, tolerant, of shade and different soil types, producing abundant seed, invading plantations (bananas, citrus, coconuts, coffee, oil palm, papaya, rubber), pastures, crops, lawns, roadsides, and wasteland. Mimosine causes thyroid problems in cattle. A fire hazard when dead.
- Small prickly shrub, up to 50 cm, creeps. Stems brown to purple, woody, branched, with sharp thorns. Leaves, hairy, alternate along stems, bearing 1-2 pairs leaf-like branches, with paired dark green leaflets. Flower stalks, with thorns, bearing bright purplish-pink, oval, flower clusters, with long stamens giving a fluffy look. Seeds, in long seed pods, splitting into 2-4, 1-seeded segments bearing bristles. Root is long and strong
- Spread: seeds, by animals, clothing; use as ornamental; moved in soil, sand, gravel.
- Biosecurity: high risk of introduction via trade as an ornamental. Among 10 worst weeds in American Samoa, French Polynesia, Guam, Solomon Islands, Tonga. On Global Invasive Species Database of alien invasive species (IUCN, 2020).
- Biocontrol: Little known.
- Cultural control: hand weed (thorns!) or hoe; mulch; vehicle hygiene; prevent overgrazing (adjust stocking rate, forcing cattle to eat young plants).
- Chemical control: in Australia: MCPA + terbutryn; fluroxypyr; dicamba; picloram + triclopyr; hexazinone + diuron. In Fiji, 2,4-D; glyphosate.

#### **Common Name**

Sensitive plant; it is also known as mimosa, common sensitive plant, or touch-me-not.

### Scientific Name

Mimosa pudica. There are a number of varieties. It is a member of the Fabaceae.

AUTHORS Grahame Jackson, Aradhana Deesh & Mani Mua

Adapted from Common sensitive plant (Mimosa pudica) (2018) Weeds of SE Qld and Northern NSW. Lucidcentral. (https://www.lucidcentral.org/editors-pick-animal-and-plant-identification-keys/key-to-weeds-of-se-qld-and-northern-nsw); and additional information from CABI (2019) Mimosa pudica (sensitive plant). Invasive Species Compendium. (https://www.cabi.org/sic/datasheet/3-4202#tosammaryOflnvasiveness); and Waterhouse DF, Norris KR (1987) Biological Control Pacific Prospects. Inkata Press, Melbourne; and from Mimosa pudica (2020) Wikipedia. Org/wiki/Mimosa\_pudica). Photos 1 &S Karan A. Rawlins, University of Georgia, Bugwood.org. Photo 4 Shaun Winterton, Aquarium and Pond Plants of the World, Edition 3, USDA APHISPPQ, Bugwood.org. Photo 7 Forest and Kim Starr, Sarr Environmental, Bugwood.org.

Produced with support from the Australian Centre for International Agricultural Research under project HORT/2016/185: Responding to emerging pest and disease threats to horticulture in the Pacific islands, implemented by the University of Queensland, in association with the Pacific Community and Koronivia Research Station, Ministry of Agriculture, Fiji.

This mini fact sheet is a part of the app Pacific Pests, Pathogens & Weeds

The mobile application is available from the Google Play Store and Apple iTunes.









Copyright © 2020. All rights reserved.