

## Asian tramp snail (369)

### Summary

- Worldwide distribution. Common in Pacific island countries.
- Shells vary, 12-16 mm diam, whitish to light brown, with or without narrow brown bands between whorls, and outward curved white lip. Hermaphrodite, capable of self-fertilization. Eggs below soil surface, hatchlings after 15 days, adults after 6 months.
- Damage direct: many crops (vegetables, trees, ornamentals) of economic importance eaten; indirect, faeces may contain rat lung worm.
- Spread on 'foot' or long distances via international plant trade.
- Biosecurity: commonly found on freight containers (hence the name 'tramp').
- Natural enemies: predatory snails and flatworms should NOT be introduced without environmental impact studies. These generalists will annihilate local snails. Use Indian runner ducks.
- Cultural control: bare earth or sand around plots; collect snails; use beer traps to attract snails.
- Chemical control: Note metaldehyde NOT encouraged: toxic to livestock, pets and humans; use baits with iron phosphate, sodium ferric EDTA - effective and safer.

### Common Name

Asian tramp snail

### Scientific Name

*Bradybaena similis*; previously, *Helix similis*.



Photo 1. Collection of Asian tramp snails, *Bradybaena similis*, showing the variation of shell colour.




Photo 2. Shells of the tramp snail, *Bradybaena similis*, showing outward curling of the lip at the mouth of the shell; this lip is white (on some shells).




Photo 4. Asian tramp snail, *Bradybaena similis*, and leaf damage on bitter gourd.



Photo 3. Asian tramp snail, *Bradybaena similis*, on bitter gourd.

AUTHOR Grahame Jackson & Mani Mua  
Information from Brodie G, Barker GM (2012) *Bradybaena similis* (Rang, 1831). Family Bradybaenidae. Introduced land snails of the Fiji Islands. Fact Sheet Series, No. 2. USP, Fiji; and Bradybaenidae. Terrestrial Mollusc Tool. USDA, University of Florida. (<http://idtools.org/id/mollusc/factsheet.php?name=Bradybaenidae>); and Asian trampsnail. Wikipedia. ([https://en.wikipedia.org/wiki/Asian\\_trampsnail](https://en.wikipedia.org/wiki/Asian_trampsnail)); and from Capinera JL, White (2014) Featured Creatures: Terrestrial snails affecting plants in Florida. UF/IFAS, University of Florida. ([http://entnemdept.ufl.edu/creatures/misc/gastro/terrestrial\\_snails.htm](http://entnemdept.ufl.edu/creatures/misc/gastro/terrestrial_snails.htm)). Photo 1 Zell H *Bradybaena similis*. ([https://commons.wikimedia.org/wiki/File:Bradybaena\\_similaris\\_01.JPG](https://commons.wikimedia.org/wiki/File:Bradybaena_similaris_01.JPG)).

Produced with support from the Australian Centre for International Agricultural Research under project HORT/2016/185: *Responding to emerging pest and disease threats to horticulture in the Pacific islands*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

Copyright © 2021. All rights reserved.



Australian Government  
Australian Centre for  
International Agricultural Research



Web edition hosted at <https://apps.lucidcentral.org/pppw>