

Celery root knot nematode (254)

Summary

- Worldwide distribution. In tropics and sub-tropics. Apart from celery, common vegetable hosts are beans, capsicum, carrot, cucumber, eggplant, ginger, lettuce, potato, tomato, and yam; common fruit crop hosts are melon, papaya, and pineapple, and many ornamentals. An important nematode disease.
- Above ground, leaves yellow with brown margins, wilt and die early; below ground, galls on roots.
- Eggs in soil hatch, young larvae enter root behind the tip, and cause cells to swell and galls to form.
- Cultural control: important - pasteurise nursery soil or use soil-less mixes; add organic matter; solarise soil for 4-6 weeks under black plastic; keep soil well-watered and use mulch; collect and burn trash after harvest; fallow land for 4-6 months; rotate with e.g., more tolerant maize, peanuts, onions, brassicas, chilli, sweetpotato, or sorghum x Sudan grass hybrids, green panic grass, or grow marigold cover crops.
- Chemical control: none recommended.

Common Name

Root knot nematode. General account of root knot nematodes (see **Fact Sheet no. 127**).

Scientific Name

Meloidogyne species. *Meloidogyne hapla*, *Meloidogyne incognita*, and *Meloidogyne javanica* all attack celery.



Photo 1. Gall on the roots of *Phaseolus* bean, caused by *Meloidogyne* species.



Photo 2. Severe infestation of root knot nematode, *Meloidogyne* species, on tomato.



Photo 4. Yellowing of celery leaves caused by root-knot nematode, *Meloidogyne* species.



Photo 3. Yellowing and browning at margins of celery leaves, caused by root-knot nematode, *Meloidogyne* species.



Photo 5. Plant in Photo 2 showing below ground symptoms: swollen roots and galls caused by root-knot nematode, *Meloidogyne* species.

AUTHOR Grahame Jackson

Information from (including Photo 2) *Diseases of vegetable crops in Australia* (2010). Editors, Denis Persley, et al. CSIRO Publishing. GrowVeg (undated) Celery root knot nematodes. (<https://www.growveg.com.au/plant-diseases/uk-and-europe/celery-root-knot-nematodes/>); and Westerdahl BB, et al. (undated) Celery Nematodes. University of California Statewide IPM Program. Agriculture and Natural Resources, University of California. (<http://ipm.ucanr.edu/PMG/r104200111.html>); and from Root-knot nematode (2017) Department of Agriculture and Fisheries. Queensland Government. Photo 1 Gerlach WWP (1988) *Plant diseases of Western Samoa*. Samoan German Crop Protection Project, Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ) GmbH, Germany.

Produced with support from the Australian Centre for International Agricultural Research under project PC/2010/090: *Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production*, implemented by the University of Queensland and the Secretariat of the Pacific Community.



Australian Government
Australian Centre for
International Agricultural Research



Web edition hosted at <https://apps.lucidcentral.org/pppw>