

## Sugarcane veneer blotch (219)

---

### Summary

- Narrow distribution. Indonesia, Oceania. On sugarcane. A minor disease.
- Spots small, oval, light green to pale yellow with thin red margins; later, surrounded by progressively larger spots, turning brown, up to 60 cm.
- Spores on the underside of leaves spread by wind.
- Cultural control: not likely to be needed.
- Chemical control: none recommended.



Photo 1. Oval spots of veneer blotch on the midrib and leaf blade of sugarcane, caused by *Deightonella papuana*. Note how the older spots enclose the younger ones.

### Common Name

Sugarcane veneer blotch

### Scientific Name

*Deightonella papuana*

AUTHOR Grahame Jackson

Photo 1 Kohler F, et al. 1997) Diseases of cultivated crops in Pacific Island countries. South Pacific Commission. Pirie Printers Pty Limited, Canberra, Australia.

Produced with support from the Australian Centre for International Agricultural Research under project PC/2010/090: Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production, implemented by the University of Queensland and the Secretariat of the Pacific Community.

Copyright © 2021. All rights reserved.



Australian Government  
Australian Centre for  
International Agricultural Research



Web edition hosted at <https://apps.lucidcentral.org/pppw>