

Lango huhukia e fua'i akau 'i Tonga (Tonga fruit flies) (171)

Fakama'opo'opo

- *Bactrocera kirki* – Lango huhukia 'o e abiu, 'avoka, kuava, mango, nonu, fekika pea 'oku toe lahi ange 'a e lango huhukia Tonga *B. facialis* ' i he famili 'e 29 'oe vesitapolo mo e 'akau fua (polo fifisi, polo melie, kuava) *B.xanthodes*- abiu,'avoka, mei 'initia, lesi, apele' initia mo ha fa'ahinga kehe.
- 'Oku fiemau 'e he lango fefine ha polotini kimu'a pea toki lava 'o fakato ha fua 'oku mo'ui.
- **Founga angamaheni hono mapule'i:** To e vesitapolo 'oku' ikai ko hono taimi totonu, kofu e fua'aki e pepa pe koe lau'i 'akau, tanaki e fua 'oku ngangana pea faka'auha, toli kimu'a te'eki 'uesia,'oku 'i ai e kalasi 'ikai sai'ia ai 'e lango.e.g koe kalasi 'o e polo fifisi moe faina 'i Fisi.
- **Founga Ngaueaki e kemikale:** ngaue'aki 'a e fana polotini tauhele lango huhukia 'oku ngaue'aki fakakomesiale e.g MPPIL, Royal Tongalure pe Bactrogel.

Common Name: Ko e kalasi lango huhukia 'e 6 'oku 'i Tonga ni. Ko e kalasi 'e 5 'oku 'i ai 'ene mahu'inga faka'ekonomika: (i) Lango huhukia Tonga (*Bactrocera facialis*) [tukukehe 'a e ongo Niua]; (ii) Lango huhukia Pasifiki (*Bactrocera xanthodes*), (iii) *Bactrocera kirki*; (iv) *Bactrocera species near passiflorae* (Ongo Niua pe); moe (v) *Bactrocera distincta*.

Scientific Name: *Bactrocera facialis*, *Bactrocera kirki*, *Bactrocera xanthodes*, *Bactrocera passiflorae*, *Bactrocera Distincta* and *Bactrocera obscura*.

Ko e lango huhukia Pasifiki, *Bactrocera xanthodes* oku vaofi 'aupito ia ki he kalasi kehe 'e 3. 'A ia ko e *Bactrocera paraxanthodea* o Niukaletonia, *Bactrocera neoxanthodes* i Vanuatu, pea moe kalasi 'oku te'eki ai ke fakahingoa 'i Ha'amoia.



Photo 1. *Bactrocera kirki*.



Photo 2. *Bactrocera facialis*.



Photo 3. *Bactrocera xanthodes*.



Photo 4. Pacific fruit fly, *Bactrocera xanthodes*, laying eggs.



Photo 5. Fruit fly eggs.



Photo 6. Fruit fly larvae or maggots.




Photo 7. Fruit fly pupae.



Photo 8. Newspaper with edges folded and stapled to form a bag in which to insert fruit to protect it from fruit flies.



Photo 9. Banana leaves used to wrap a fruit bunch to prevent attack by fruit flies, other pests, and to promote uniform ripening, in Papua New Guinea.

Copyright © 2023. All rights reserved.



Australian Government
Australian Centre for
International Agricultural Research



Web edition hosted at <https://apps.lucidcentral.org/pppw>