

Cryptocarya densiflora Blume

Family:
Lauraceae

Blume, C.L. (1826) *Bijdragen tot de Flora van Nederlandsch Indie* 11: 556. Type: *Crescit in sylvis obscuris in montibus Saleyer*

Common name:
Cinnamon Laurel; Laurel, Cinnomon; Laurel, White; White Laurel

Stem

Sapwood surface corrugated. A thin pale brown layer generally visible beneath the subrhynchidome layer before the first section of the outer blaze. Often a poorly formed tree with a number of coppice shoots at the base.

Leaves

Twigs fluted, clothed in straight brown, appressed hairs when young, but sometimes almost glabrous when older. Leaf blades about 8-15 x 2.5-6 cm, glaucous on the underside, clothed in straight, pale brown, appressed hairs when young but eventually becoming almost glabrous. Small foveoles sometimes visible in the axils of the basal pair of lateral veins. Midrib depressed, occasionally flush with the upper surface. Petioles channelled on the upper surface. Oil dots visible with a lens.

Flowers

Inflorescence panicle, sometimes exceeding the leaves. Flowers without any obvious perfume. Lower half of the perianth tube pubescent on the inner surface. Tepals about 0.9-2.7 mm long, pubescent on the outer surface. Ovary and style glabrous.

Fruit

Fruits longitudinally ribbed, depressed globular, about 11-14 x 13-19.5 mm. Cotyledons white or cream.

Seedlings

First pair of leaves ovate to elliptic, about 23-40 x 14-25 mm, glaucous on the underside. At the tenth leaf stage: leaf blade glaucous on the underside, upper surface of leaves and petioles glabrous; oil dots very small, visible only with a lens. Seed germination time 110 to 269 days.

Distribution and Ecology

Occurs in CYP, NEQ and CEQ. Altitudinal range from 450-1200 m. Grows as an understory tree in well developed upland and mountain rain forest. Also occurs in Malesia.

Natural History & Notes

This species has little commercial value as it seldom grows large enough to produce millable logs and the peculiar structure of the wood does not encourage use of the timber. Wood specific gravity 0.70 -0.72. Hyland (1989).

Synonyms

Cryptocarya cinnamomifolia Benth., *Flora Australiensis* 5: 298(1870), Type: Queensland. Rockingham Bay, Dallachy.

RFK Code

5



Flowers [not vouchered]. © G.

Sankowsky




Fruit, several views, cross and longitudinal sections. © W. T. Cooper

Cooper



Habit, leaves and flowers. ©

CSIRO



Scale bar 10mm. © CSIRO



Cotyledon stage, hypogeal germination. © CSIRO



10th leaf stage. © CSIRO



Web edition hosted at <https://apps.lucidcentral.org/rainforest>