

WATTLE

Acacias of Australia

Acacia lanigera var. gracilipes Benth.



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.



Acacia lanigera var. gracilipes occurrence map.
Occurrence map generated via Atlas of Living
Australia ([https://www.ala.org.au](http://www.ala.org.au)).

Family

Fabaceae

Distribution

Occurs in a restricted area from near Mountain Ck (S of Mt Deddick) and the Genoa R. area in eastern Vic. to the neighbouring Wallagaraugh R. area in N.S.W.

Description

Branchlets densely tomentulose with patent to subappressed hairs. Phyllodes commonly oblong-elliptic and straight, (1.5–) 3–6 cm long, (3–) 4–9 mm wide, with few anastomoses; gland 0–1 mm above pulvinus, not prominent, commonly obscured by indumentum that extends from pulvinus. Peduncles 5–12 mm long, glabrous; flower-heads c. 5 mm diam.

Habitat

Usually grows among granite boulders in open forest or shrubland.

Specimens

N.S.W.: junction of Imlay Ck and Wallagaraugh R., D.E.Albrecht 3938 (MEL). Vic.: East Gippsland: Genoa R., c. 1 km downstream from the Tasker track crossing, Wangarabell area, D.E.Albrecht 4878 (MEL, NSW); Deddick Track, 1.2 km NW of Mt Joan, 13.8 km S Mt Deddick, S.J.Forbes 162 (MEL, NSW, PERTH).


FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

R.S.Cowan, B.R.Maslin

This identification key and fact sheets are available as a mobile application:



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.